



EVO-MPMI: From fundamental science to practical applications

Sebastian Schornack¹ and Sophien Kamoun²

Abstract

In the unending coevolutionary dance between plants and microbes, each player impacts the evolution of the other. Here, we provide an overview of the burgeoning field of evolutionary molecular plant–microbe interactions (EVO-MPMI)—the study of mechanisms of plant–microbe interactions in the context of their evolutionary history—tracing its progression from foundational science to practical implementation. We present a snapshot of current research and delve into central concepts, such as conserved features and convergent evolution, as well as methodologies such as ancestral reconstruction. Moreover, we shed light on the practical applications of EVO-MPMI, particularly within the realm of disease control. Looking ahead, we discuss potential future trajectories for EVO-MPMI research, spotlighting the innovative tools and technologies propelling the discipline forward.

Addresses

¹ Sainsbury Laboratory, University of Cambridge, Cambridge, United Kingdom

² The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, United Kingdom

Corresponding author: Kamoun, Sophien (Sophien.Kamoun@tsl.ac.uk)

Current Opinion in Plant Biology 2023, 76:102469

This review comes from a themed issue on **Biotic interactions 2024**

Edited by **Pierre-Marc Delaux** and **Jacqueline Monaghan**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online xxx

<https://doi.org/10.1016/j.pbi.2023.102469>

1369-5266/© 2023 Elsevier Ltd. All rights reserved.

Une valse à mille temps: The coevolutionary waltz between plants and microbes

While plant hosts are known to shape the evolution of symbionts and pathogens by driving changes within microbial genomes, the microbes leave their own mark on the evolution of their plant hosts. This has resulted in specific genetic signatures in microbial and plant genomes that are associated with past and ongoing interactions. These footprints provide evidence of the dynamic and reciprocal nature of plant–microbe coevolution, and can provide new insights into the underlying molecular mechanisms that drive these interactions. This

is the basis of the emerging field of evolutionary molecular plant–microbe interactions (EVO-MPMI)—the study of mechanisms of plant–microbe interactions in the context of their evolutionary history. Using this integrated multi-disciplinary approach, researchers can gain new insights into the complex and dynamic processes that shape the diversity and adaptability of these organisms. These coevolutionary systems are exceptional in that they can be studied over thousands of tempos, from millions of years to very short timescales, revealing anything from macroevolutionary patterns to rapid evolutionary adaptations [1,2].

Here we survey the EVO-MPMI field, covering recent developments from fundamental science to practical applications. The article explores key concepts and methods, and current state of research. It will also highlight the practical applications of EVO-MPMI in the areas of disease control. In addition, we discuss the future directions of EVO-MPMI research, including the emerging tools and technologies that are driving the field forward.

Beyond model species: The hidden diversity of molecular mechanisms

Our understanding of molecular mechanisms of plant–microbe interactions tends to be limited to a few model species and does not do justice to the huge biodiversity of interacting organisms. There is a whole world of microbial and plant diversity out there but not all of it is as accessible as some model species. Investigating molecular mechanisms in understudied, and experimentally challenging systems has revealed novel pathways and unexpected innovations. For example, the recently compiled RefPlantNLR database underscores that, out of 59 orders of flowering plants, the intracellular immune receptors of the nucleotide-binding leucine-rich repeat (NLR) class have been experimentally validated in only 11 orders [3].

Recent years have seen a surge in research in non-flowering and non-vascular plants. *Marchantia polymorpha* was particularly quickly adopted as a popular experimental system [4]. Its mostly streamlined genome without whole genome duplications allows to test functional conservation of genes across large evolutionary distances. Like all plants the *Marchantia*

genome encodes for conserved and lineage specific molecular mechanisms. Conserved biochemical defenses including the induction of pathogenesis-related proteins and phenylpropanoid biosynthesis enzymes underpin *Marchantia* responses to oomycete infection. However, the purple pigment which accumulates at infection sites is not an anthocyanin, but a combination of chemically different and liverwort lineage-specific auronidins [5,6]. Bryophyte genome sequencing also uncovered new types of NLR genes with uncommon N-termini coding for putative hydrolases and kinases (for a recent overview see the study by Chia et al. [7]). If truly lineage specific, these domains might aid in the engineering of crop immune receptors against angiosperm specific pathogens, which might not have evolved mechanisms to inhibit them.

Diversity in plant–microbe interactions is also evident in nitrogen-fixing root nodules. Plant-secreted nodule-specific cysteine-rich (NCR) peptides probably evolved from antimicrobial, short, cysteine-rich defensin-like peptides found in fungi, plants and animals. NCR peptides regulate the symbiosis of nitrogen-fixing rhizobacteria by controlling the terminal differentiation of the microsymbiont (for a review see the study by Lima et al. [8]). NCR peptides have evolved independently in two clades of legumes: the Inverted-repeat-lacking (IRLC, eg. *Medicago truncatula*) and the Dalbergoid clade (eg. *Aeschynomene indica*). Other legumes, such as soybean and *Lotus japonicus*, do not have NCR peptides. Likely, other mechanisms control the microsymbiont in these plants. Recent surprising results revealed that the nodule restricted expression of NCRs is not controlled by a lineage-specific mechanism. Instead, acquisition of NCR genes also involved recruitment of nodule-specific promoters which are regulated by existing transcription factors [9]. Similar to defensins, NCR peptides have antimicrobial activity and can terminate bacterial cell divisions with promising leads as antibiotics [8].

Microbe-encoded, host-translocated proteins called effectors target plant processes and are central to a microbe's success in colonising a plant. Functional effector studies based on the expression of effectors directly in plants have significantly added to our understanding of their biology. For Phytoplasmas, bacteria of the class of Mollicutes which reside within plant phloem cells and are transmitted by sap-sucking insects this was a real game changer. Phytoplasma effector expression in plants has led to the same morphological phenotypes observed during phytoplasma infections [10–12]. Then, the identification of host targets of these effectors was a key step in understanding underlying mechanisms such as SAP54's ability to degrade transcription factors in an ubiquitination-independent manner [12]. Mechanisms like these have wide potential for application beyond plants and highlight

the diversity of molecular mechanisms that occur beyond the classic model pathogen species.

Recently, AlphaFold2 and similar deep-learning algorithms have enabled structure-model based comparisons in pathogens as well as genetically challenging arbuscular mycorrhiza fungi [13,14] and led to the identification of structurally similar effector candidates that are not conserved at the primary sequence level [15] (see also below the section on convergent evolution). Facing this wealth of candidate effectors we now need to understand the molecular mechanisms underpinning their functions [16]. This requires the development of new assays and readouts beyond testing suppression of immune pathways and assessing their subcellular distribution *in planta*. Knowing how structure confers function is also a starting point for protein engineering towards cell process control in disease and developmental contexts.


Conserved genes enable a wider perspective on structure and function

Comparative sequence analyses, conducted within a robust evolutionary framework, have emerged as a powerful approach to unravel molecular mechanisms and generate experimentally testable hypotheses. We illustrate this concept with examples centered on NLR class of plant immune receptors (Figure 1).

Recent studies have employed a phylogenomic approach to investigate the molecular evolution of the NLR protein ZAR1 in flowering plants (angiosperms) [17–19]. These investigations unveiled over 100 ZAR1 orthologues spanning monocots, magnoliids, and eudicots, indicating the exceptionally conserved nature of ZAR1, tracing its origins back to the early angiosperm lineages in the Jurassic period, around 220 to 150 million years ago. ZAR1 is known for indirectly detecting bacterial pathogens through binding pseudokinase decoys of the RLCK family. Remarkably, Adachi et al. [19] demonstrated that ZAR1 has been partnering with RLCKs for more than 150 million years, as supported by functional reconstructions of ZAR1-RLCK pairs from distantly related plant species (see Figure 1). These RLCKs are found in gene clusters across diverse angiosperms and are believed to have evolved from receptor kinases (RKs) involved in the recognition of microbial patterns. These findings suggest the presence of RK pathways dedicated to immunity during the Jurassic period, and that these pathways were already targeted by pathogens. Notably, the conserved ZAR1 kinase binding interface within the leucine-rich repeats supports this model, highlighting the enduring significance of this interaction across ZAR1 orthologues.

A striking conserved feature of ZAR1 is the MADA $\alpha 1$ helix, which plays a crucial role in defining the anion channel of the activated oligomeric resistosome [20,21]. This MADA motif is functionally conserved across a

Figure 1



Conserved genes: A gateway to molecular mechanisms. (a) Workflow to identify conserved protein motifs and interfaces. This adapted workflow combines phylogenomics analysis, sequence conservation, and 3D structure comparisons. By generating experimentally testable hypotheses, this pipeline can help unveil novel biochemical features, offering insights into their functional significance. The figure is adapted from the study by Adachi et al. [19]. (b) Conserved sequence patterns in the NLR immune receptor ZAR1 based on comparative analyses of an orthologue series. The schematic representation of ZAR1 protein highlights the position of these conserved sequence patterns across ZAR1 orthologues. Residues that have been functionally validated in Arabidopsis ZAR1 are marked with red asterisks. Motifs were identified using MEME and conservation scores calculated with ConSurf. The figure is adapted from Adachi et al. [17,19].

diverse range of angiosperms. In a recent study, Chia et al. [7] demonstrated that the MADA motif can be traced back to non-flowering plants, suggesting its origin to be at least 450 million years ago. Interestingly, in certain lineages of non-flowering plants, a variant of the MADA motif known as MAEPL defines the $\alpha 1$ helix of CC-NLRs (coiled-coil nucleotide-binding leucine-rich repeat receptors). Remarkably, this MAEPL variant is functionally interchangeable with the angiosperm MADA N-terminus [7]. However, whether the occurrence of the MAEPL variant is a consequence of historical contingency or represents an adaptive feature remains an intriguing question that requires further investigation.

This study exemplifies the value of integrating sequence conservation data with 3D structural analyses, as depicted in Figure 1. Looking ahead, as more protein structures become available, either through experimental efforts or computational predictions, the availability of orthologue series for a given protein will prove

instrumental in uncovering conserved features and generating hypotheses regarding biochemical activities. Focusing solely on a single protein, such as the canonical Arabidopsis ZAR1, without tapping into the wealth of available sequence diversity, is unlikely to yield the same depth of insights. Additionally, variable features that deviate from the overall conserved sequences within an orthologue series can also provide valuable hypotheses. For instance, in eudicots, ZAR1 duplicated into the paralogous ZAR1-SUB proteins, which lack the kinase binding interface [17–19]. This observation led Adachi et al. [19] to postulate that ZAR1-SUB proteins may have neo-functionalised to bind ligands other than RLCKs, offering a captivating avenue for further exploration and investigation.

Journeying through evolution: Ancestral reconstructions inform mechanistic models

Ancestral reconstructions of extinct proteins at pivotal phylogenetic nodes have gained popularity in EVO-


MPMI studies [22–26] (Figure 2). These reconstructions play a crucial role in unraveling the evolutionary pathways taken by a protein, shedding light on evolutionary transitions and adaptive characteristics that provide valuable insights into molecular mechanisms. This approach allows researchers to reconstruct ancestral protein sequences that are no longer present in extant organisms. By tracing back evolutionary history, these reconstructions provide a glimpse into the ancestral forms of proteins, offering a deeper understanding of their functional and structural changes over time.

In a study on the neo-functionalisation of a *Phytophthora infestans* protease inhibitor effector EPIC1 after a host jump, the ancestral gene was resurrected to investigate the hypothesis regarding shifts in its activity spectrum over evolutionary time [23]. The inhibition spectrum of the resurrected protein indicated that the ancestral protein had distinct activities from the derived protein from the sister species *Phytophthora mirabilis* consistent with the hypothesis that this effector has adapted to protease targets unique to its host plant *Mirabilis jalapa* [23].

Ancestral reconstructions enabled a plausible model of coevolution between APiKL effectors from the blast fungus pathogen *Magnaporthe oryzae* and its host grass plants [24] (Figure 2). Ancestral reconstructions established the directionality of evolution and determined that the ancestral form of the APiKL2 effector, D66 (GAT codon), preceded the derived form, N66 (AAT codon). Based on this insight, the D66N polymorphism is more likely to reflect an expansion in host-target binding rather than evasion of host immunity, and thus was proposed to be an adaptive feature that enabled binding to a new host target [24].

Ancestral reconstructions revealed the evolutionary history of rice immune receptors, Pik-1 and Pik-2, and their integrated HMA domain [24] (Figure 2). By reconstructing the evolutionary history of Pik-1 and Pik-2, coupled with functional studies, Bialas et al. [25] tested hypotheses regarding the adaptive evolution of the HMA domain. Employing ancestral sequence reconstruction, they demonstrated that two allelic variants of Pik-1 independently evolved from a weakly binding ancestral state to exhibit high-affinity binding to the blast fungus effector AVR-PikD. In another study,

Figure 2



Ancestral reconstructions inform mechanistic models. (a) The Asp-66-Asn polymorphism in the APiKL2 effector of the blast fungus *Magnaporthe oryzae* resulted in expanded binding to target small heavy metal-associated (sHMA) proteins. The figure depicts the host-specific lineages of *M. oryzae*, with annotation of the Asp-66/Asn-66 polymorphism (blue for Asp-66 and red for Asn-66). The model indicates that the Asn-66 polymorphism emerged prior to the differentiation of host-specialised lineages, and is likely to correspond to an expansion of the binding spectrum to sHMA proteins rather than evasion of detection by an immune receptor. The Figure is adapted from the study by Bentham et al. [24]. (b) Molecular convergence of rice immune receptors Pikp-1 and Pikm-1 towards high affinity for the blast fungus effector AVR-PikD. Ancestral reconstructions revealed that the HMA domains of the Pikp-1 and Pikm-1 receptors have undergone convergent evolution following distinct evolutionary paths. The Pikp-HMA domain evolved through AV-VE adaptations in the IAQVV/LVKIE region, while the Pikm-HMA domain acquired ANK-VKE mutations in the MKANK/EMVKE region. Bialas et al. [25] proposed a model in which the high-affinity binding of Pikp-1 and Pikm-1 allelic variants to the AVR-PikD effector has occurred relatively recently during the evolutionary history of this immune receptor. The Figure is adapted from the study by Bialas et al. [25].

Snoeck et al. [26] used a combination of phylogenomics and ancestral reconstructions to show that INR, a cell-surface immune receptor that responds to insectin molecules from caterpillars, emerged around 28 million years ago as a result of incremental mutations in the ancient form of the receptor [26].

Improbable destinities: Insights from convergent evolution

Convergent evolution occurs when different species or groups of organisms evolve similar adaptations to similar environmental cues [27]. Thus, studying examples of convergence can provide evidence for natural selection and adaptation and reveal advantageous traits.

An example are plant-pathogen effectors that converge on a limited number of host pathways, suggesting that there are critical processes for successful infection that pathogens have evolved to manipulate. Phylogenetically and structurally different effectors from unrelated microbial species target conserved signaling pathways and regulatory proteins important for host defense, such as immune signaling pathways [28,29], or the cellular endomembrane trafficking pathway [30]. Convergence can reveal the constraints and options underpinning a specific evolutionary process. Molecular studies are important to resolve whether effectors of divergent microbes target identical host proteins (Reviewed here: [31]) or different proteins within the same pathway or use the same molecular mechanism to do so. An example for the latter are bacterial and oomycete pathogens which independently have evolved effectors that act as substrate mimics for host 14-3-3 proteins [32,33]. Understanding the mechanisms by which effectors manipulate key pathways and processes provides opportunities for the development of new strategies for disease control.

Genes underpinning plant immunity also display signs of convergent evolution. Immune receptors underpin distinct elicitor recognition in related Solanaceous plants [34]. Barley and Arabidopsis have convergently evolved different nucleotide-binding leucine-rich repeat (NLR) proteins to detect effectors that structurally modify PBS1-like kinases [35].

Convergence also aids in reconstructing evolutionary history. Genome sequence analyses have shown the independent evolution of the ectomycorrhizal lifestyle of several independent fungal lineages from their saprotrophic ancestors. This included the convergent losses of different plant cell wall-degrading enzymes [36]. In the endomycorrhiza formed between most land plants and arbuscular mycorrhiza (AM) fungi, PHT1 type phosphate transporters play a key role in nutrient uptake into plant cells. While PT4/PT11 clade PHT1 phosphate transporters are transcriptionally upregulated

in most studied angiosperm models, some Solanaceae and the bryophyte *Lunularia cruciata* upregulate PHT1-type transporters from non-PT4 clades, suggesting that the molecular mechanisms underpinning nutrient transfer may have evolved convergently [37–39].


Evolutionary tinkering paves the way for bioengineering

In the words of the eminent biologist Francois Jacob, evolution can be likened to a skilled tinkerer, subtly modifying biological molecules over the course of millions of years. Biotechnologists have taken inspiration from this natural process, striving to replicate and expedite evolution within the confines of our laboratories. In the realm of EVO-MPMI, numerous examples demonstrate the transformative power of evolutionary reconstructions in guiding bioengineering endeavors.

The evolutionary trajectory of TAL effectors (TALEs), predominantly found in the bacterial pathogens and symbionts *Xanthomonas* and *Burkholderia*, has captivated researchers in the field. These effectors have evolved the remarkable ability to bind DNA that is governed by a protein-to-DNA cipher that enables TALEs to bind specific DNA sequences [40–42] (Figure 3). The initial discovery of the cipher was based on comparisons of related and unrelated TALEs converging onto the same promoter elements. Subsequently, evolutionary studies have provided insights into how TALEs evolve to maintain binding to susceptibility gene promoters [43]. This knowledge has guided the design and engineering of synthetic TALE-derived DNA-binding proteins, which can be combined with other domains to enable gene editing and modulation of transcriptional activities, among many applications [44]. TALEs have also applications in plant disease resistance (Figure 3). Among these is the identification and design of decoy sequences that trap pathogen TALEs to activate executors of immunity [45] and the use of experimental evolution to predict TALE adaptations that evade binding to sequences that activate immunity [46].

Another area where evolutionary reconstructions have been instrumental is in the realm of NLR-IDs (NLR immune receptors with integrated domains). This remarkable process—when an unconventional domain is integrated into a classical NLR architecture to bait pathogens—has occurred hundreds of times throughout plant NLR evolution [47,48]. By reconstructing the evolutionary history of the Pik NLR pair, which includes an NLR-ID [49,50], Bialas et al. [24] and De la Concepcion et al. [51] unlocked the bioengineering principles for building made-to-order NLR-ID proteins with enhanced functionalities. This work has set the stage for pikobodies, an approach that seeks to mimic the evolutionary process by integrating animal antibody

Figure 3



Evolutionary tinkering of TAL effectors guide bioengineering and disease resistance strategies. (a) Structure of TALE proteins. TALE proteins consist of a series of repeats, each containing two hypervariable amino acids that bind to DNA following a specific code. T3SS: Type III secretion signal, NLS: nuclear localisation signal, AAD: acidic transcription activation domain. 12/13th repeat-variable diresidues RVD are listed in each repeat. The code table lists common RVD-to-DNA base specificities. (b) Workflow for the identification and design of sequences that trap TALEs to activate executors of immunity. The pipeline consists of using knowledge on TAL effectors, to identify susceptibility genes and their loss-of-function alleles in plants targeted by specific *Xanthomonas* species. The TAL effector repertoires are determined from strains collected through field surveys. To identify the corresponding susceptibility genes, three strategies are employed: assessing the TAL effector's virulence role in plants, using bioinformatics tools to predict TAL effector targets in the host genome, and using RNA profiling to spot TAL effector-dependent genes. After identifying potential genes, their functional roles are examined, leading to the discovery of key susceptibility hubs. Natural or edited genetic variations in these genes can then offer new avenues for plant disease resistance. Adapted from the study by Hutin et al. [45]. (c) Experimental evolution pipeline of TALE proteins. The figure illustrates the experimental evolution of TALEs randomised in their DNA binding sequences and subjected to cycles of infection on resistant plants. Following 30 cycles of experimental evolution, the recovered TALEs were transformed into *Escherichia coli* for sequencing to identify adaptations that evade binding to sequences that activate immunity. Adapted from the study by Teper et al. [46].

domains into NLR proteins [52]. If nature figures out how to bioengineer an NLR-ID every few millions of years, why can't we make this routine in our laboratories?

Evolutionary reconstructions have also helped produce a framework for how functionally linked immune receptors undergo co-adaptation to provide an effective and regulated immune response against pathogens [48,51]. This has revealed, for instance, how mismatched Pik-1/Pik-2 pairs can lead to autoimmune or hypoactive phenotypes and which amino acids modulate these activities [51], helping to add to the toolbox for pikobody bioengineering [52,53].

Future prospects

The future of EVO-MPMI appears promising, with opportunities for both scientific advancements and practical applications. The development of innovative pipelines and methodologies will be crucial. Such advancements

will enable comprehensive and efficient analysis of large-scale data, incorporating high-throughput sequencing, comparative genomics, and functional assays. By taking a holistic view of the complex interplay between plants and microbes, scientists will be able to interpret the underlying molecular mechanisms driving these interactions in the light of evolution.

One key avenue that is set to shape the field's trajectory is the integration of molecular evolution with structural biology (Figures 1–2). By combining these two powerful approaches, researchers can delve into the intricate mechanisms and adaptive features of plant-microbe interactions at the molecular level, unraveling their evolutionary dynamics and gaining deeper insights.

Nobel Prize laureate Frances H. Arnold's groundbreaking concept of 'innovation by evolution' [54] serves as an inspiration for EVO-MPMI researchers.

Drawing upon the power of natural selection, they can leverage evolutionary principles to drive practical applications. By identifying and optimising proteins and molecules with desired functionalities, the field holds the potential for groundbreaking biotechnological applications.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Sophien Kamoun has patent Plant disease resistance genes pending to The Sainsbury Laboratory. S.S. holds patents on TALEs. S.K. receives funding from industry and has filed patents on NLR biology.

Data availability

No data was used for the research described in the article.

Acknowledgements

We thank past and present members of our laboratories as well as several colleagues and three anonymous reviewers for numerous discussions and ideas. This article is not intended as a comprehensive review of the topic; consequently, several excellent studies may not have been included. The text was written and edited with the assistance of ChatGPT. Figures were adapted as indicated in the legends from articles published under CC-BY license. Sebastian receives funding primarily from the Gatsby Charitable Foundation, The Royal Society, Biotechnology and Biological Sciences Research Council (BBSRC, UK) and European Research Council (ERC). Sophien receives funding primarily from the Gatsby Charitable Foundation, BBSRC, UK, and ERC.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Upton JL, Zess EK, Białas A, Wu CH, Kamoun S: **The coming of age of EvoMPMI: evolutionary molecular plant-microbe interactions across multiple timescales.** *Curr Opin Plant Biol* 2018, **44**:108–116, <https://doi.org/10.1016/j.pbi.2018.03.003>.
2. Delaux PM, Schornack S: **Plant evolution driven by interactions with symbiotic and pathogenic microbes.** *Science* 2021, **371**, eaba6605, <https://doi.org/10.1126/science.aba6605>.
3. Kourelis J, Sakai T, Adachi H, Kamoun S: **RefPlantNLR is a comprehensive collection of experimentally validated plant disease resistance proteins from the NLR family.** *PLoS Biol* 2021, **19**, e3001124, <https://doi.org/10.1371/journal.pbio.3001124>.
4. Bowman JL, Arteaga-Vazquez M, Berger F, Briginshaw LN, Carella P, Aguilar-Cruz A, Davies KM, Dierschke T, Dolan L, Dorantes-Acosta AE, Fisher TJ, Flores-Sandoval E, Futagami K, Ishizaki K, Jibrán R, Kanazawa T, Kato H, Kohchi T, Levins J, Lin SS, Nakagami H, Nishihama R, Romani F, Schornack S, Tanizawa Y, Tsuzuki M, Ueda T, Watanabe Y, Yamato KT, Zachgo S: **The renaissance and enlightenment of Marchantia as a model system.** *Plant Cell* 2022, **34**:3512–3542, <https://doi.org/10.1093/plcell/koac219>.
5. Berland H, Albert NW, Stavland A, Jordheim M, McGhie TK, Zhou Y, Zhang H, Deroles SC, Schwinn KE, Jordan BR, Davies KM, Andersen ØM: **Auronidins are a previously unreported class of flavonoid pigments that challenges when anthocyanin biosynthesis evolved in plants.** *Proc Natl Acad Sci USA* 2019, **116**:20232–20239, <https://doi.org/10.1073/pnas.1912741116>.
6. Davies KM, *et al.*: **Evolution and function of red pigmentation in land plants.** *Ann Bot* 2022, **130**:613–636, <https://doi.org/10.1093/aob/mcac109>.
7. Chia KS, Kourelis J, Vickers M, Sakai T, Kamoun S, Carella P: **The N-terminal executioner domains of NLR immune receptors are functionally conserved across major plant lineages.** *bioRxiv* 2022, <https://doi.org/10.1101/2022.10.19.512840>.
This study traces back the emergence of the major classes of plant NLRs to non-flowering lineages and experimentally validates the immune activation function of representatives of these lineages.
8. Lima RM, Kylarová S, Mergaert P, Kondorosi É: **Unexplored arsenals of legume peptides with potential for their applications in medicine and agriculture.** *Front Microbiol* 2020, **18**:1307, <https://doi.org/10.3389/fmicb.2020.01307>.
9. Zhang S, Wang T, Lima RM, *et al.*: **Widely conserved AHL transcription factors are essential for NCR gene expression and nodule development in Medicago.** *Nat Plants* 2023, **9**:280–288, <https://doi.org/10.1038/s41477-022-01326-4>.
10. Sugio A, Kingdom HN, MacLean AM, Grieve VM, Hogenhout SA: **Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating plant development and defense hormone biosynthesis.** *Proceedings of the National Academy of Sciences USA* 2011, **108**:E1254–E1263, <https://doi.org/10.1073/pnas.1105664108>.
11. MacLean AM, Orlovskis Z, Kowitzanich K, Zdziarska AM, Angenent GC, Immink RG, Hogenhout SA: **Phytoplasma effector SAP54 hijacks plant reproduction by degrading MADS-box proteins and promotes insect colonization in a RAD23-dependent manner.** *PLoS Biol* 2014, **12**, e1001835, <https://doi.org/10.1371/journal.pbio.1001835>.
12. Huang W, MacLean AM, Sugio A, Maqbool A, Busscher M, Cho ST, Kamoun S, Kuo CH, Immink RG, Hogenhout SA: **Parasitic modulation of host development by ubiquitin-independent protein degradation.** *Cell* 2021, **184**:5201–5214.e12, <https://doi.org/10.1016/j.cell.2021.08.029>.
13. Seong K, Krasileva KV: **Prediction of effector protein structures from fungal phytopathogens enables evolutionary analyses.** *Nature Microbiology* 2023, **8**:174–187, <https://doi.org/10.1038/s41564-022-01287-6>.
This paper is a glimpse into the future, demonstrating that computational predictions of protein structures can reveal previously overlooked similarities between pathogen effector proteins.
14. Derbyshire MC, Raffaele S: **Surface frustration re-patterning underlies the structural landscape and evolvability of fungal orphan candidate effectors.** *bioRxiv* 2023, <https://doi.org/10.1101/2023.01.06.522876>.
15. Teulet A, Quan C, Evangelisti E, Wanke A, Yang W, Schornack S: **A pathogen effector FOLD diversified in symbiotic fungi.** *New Phytol* 2023, <https://doi.org/10.1111/nph.18996>.
Another study exploiting AlphaFold2 to reveal unexpected similarities between effector proteins, in this case from symbionts and pathogens.
16. Homma F, Huang J, van der Hoorn RAL: **Alphafold-multimer predicts cross-kingdom interactions at the plant-pathogen interface.** *bioRxiv* 2023, <https://doi.org/10.1101/2023.04.03.535425>.
This study shows that AI-based predictions of protein–protein interactions can reveal hidden activities of pathogen effectors.
17. Adachi H, Sakai T, Kourelis J, Pai H, Hernandez JLG, Maqbool A, Kamoun S: **Jurassic NLR: conserved and dynamic evolutionary features of the atypically ancient immune receptor ZAR1.** *bioRxiv* 2020, <https://doi.org/10.1101/2020.10.12.333484>.
An example of how comparative studies of an ortholog series can yield insights into the structure and function of a protein beyond the study of a representative from a single species.
18. Gong Z, Qi J, Hu M, Bi G, Zhou JM, Han GZ: **The origin and evolution of a plant resistosome.** *Plant Cell* 2022, **34**:1600–1620, <https://doi.org/10.1093/plcell/koac053>.
19. Adachi H, Sakai T, *et al.*: **Jurassic NLR: conserved and dynamic evolutionary features of the atypically ancient immune receptor ZAR1.** *Plant Cell* 2023, [koad175](https://doi.org/10.1093/plcell/koac175).
20. Wang J, Hu M, Wang J, Qi J, Han Z, Wang G, Qi Y, Wang H-W, Zhou J-M, Chai J: **Reconstitution and structure of a plant NLR**

- resistosome conferring immunity.** *Science* 2019, **364**, eaav5870, <https://doi.org/10.1126/science.aav5870>.
21. Adachi H, Contreras MP, Harant A, Wu CH, Derevnina L, Sakai T, Duggan C, Moratto E, Bozkurt TO, Maqbool A, Win J, Kamoun S: **An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species.** *eLife* 2019, **8**, e49956, <https://doi.org/10.7554/eLife.49956>.
 22. Thornton J: **Resurrecting ancient genes: experimental analysis of extinct molecules.** *Nat Rev Genet* 2004, **5**:366–375.
 23. Dong S, Stam R, Cano LM, Song J, Sklenar J, Yoshida K, Bozkurt TO, Oliva R, Liu Z, Tian M, Win J, Banfield MJ, Jones AME, van der Hoorn RAL, Kamoun S: **Effector specialization in a lineage of the Irish potato famine pathogen.** *Science* 2014, **343**:552–555.
 24. Bentham AR, Petit-Houdenet Y, Win J, Chuma I, Terauchi R, Banfield MJ, Kamoun S, Langner T: **A single amino acid polymorphism in a conserved effector of the multihost blast fungus pathogen expands host-target binding spectrum.** *PLoS Pathog* 2021, **17**, e1009957.
This study illustrates how ancestral reconstructions can distinguish between different hypotheses about the activities of a pathogen effector.
 25. Bialas A, Langner T, Harant A, Contreras MP, Stevenson CEM, Lawson DM, Sklenar J, Kellner R, Moscou MJ, Terauchi R, Banfield MJ, Kamoun S: **Two NLR immune receptors acquired high-affinity binding to a fungal effector through convergent evolution of their integrated domain.** *eLife* 2021, **10**, e66961.
A comprehensive study that combines phylogenomics and biochemical analyses to reconstruct the evolution of an NLR immune receptor building on ancestral reconstructions.
 26. Snoeck S, Abramson BW, Garcia AGK, Egan AN, Michael TP, Steinbrenner AD: **Evolutionary gain and loss of a plant pattern-recognition receptor for HAMP recognition.** *eLife* 2022, **11**, e81050, <https://doi.org/10.7554/eLife.81050>.
Another study that uses ancestral reconstructions to trace back the emergence of a cell surface immune receptor and the evolutionary transition to the modern form.
 27. Losos JB, Peterson M: *Improbable destinies: fate, chance, and the future of evolution.* New York: Riverhead Books; 2017.
 28. Song J, Win J, Tian M, Schornack S, Kaschani F, Ilyas M, van der Hoorn RAL, Kamoun S: **Two effectors secreted by unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3.** *Proc Natl Acad Sci USA* 2009, **106**:1654–1659.
 29. Derevnina L, Contreras MP, Adachi H, Upson J, Vergara Cruces A, Xie R, Sklenar J, Menke FLH, Mugford ST, MacLean D, Ma W, Hogenhout S, Govere A, Maqbool A, Wu CH, Kamoun S: **Plant pathogens convergently evolved to counteract redundant nodes of an NLR immune receptor network.** *PLoS Biol* 2021, **19**, e3001136.
 30. Petre B, Contreras MP, Bozkurt TO, Schattat MH, Sklenar J, Schornack S, Abd-El-Hallem A, Castells-Graells R, Lozano-Duran R, Dagdas YF, Menke FLH, Jones AME, Vossen JH, Robatzek S, Kamoun S, Win J: **Host-interactor screens of *Phytophthora infestans* RXLR proteins reveal vesicle trafficking as a major effector-targeted process.** *Plant Cell* 2021, **33**:1447–1471.
 31. Carella P, Evangelisti E, Schornack S: **Sticking to it: phyto-pathogen effector molecules may converge on evolutionarily conserved host targets in green plants.** *Curr Opin Plant Biol* 2018, **44**:175–180.
 32. Evangelisti E, Guyon A, Shenhav L, Schornack S: **FIRE mimics a 14-3-3-binding motif to promote *Phytophthora palmivora* infection.** *Mol Plant Microbe Interact* 2023 PMID, 36734982.
 33. Teper D, Salomon D, Sunitha S, Kim JG, Mudgett MB, Sessa G: **Xanthomonas euvesicatoria type III effector XopQ interacts with tomato and pepper 14-3-3 isoforms to suppress effector-triggered immunity.** *Plant J* 2014, **77**:297–309.
 34. Chen Z, Liu F, Zeng M, Wang L, Liu H, Sun Y, Wang L, Zhang Z, Chen Z, Xu Y, Zhang M, Xia Y, Ye W, Dong S, Govers F, Wang Y, Wang Y: **Convergent evolution of immune receptors underpins distinct elicitor recognition in closely related Solanaceous plants.** *Plant Cell* 2023, **35**:1186–1201.
 35. Carter ME, Helm M, Chapman AVE, Wan E, Restrepo Sierra AM, Innes RW, Bogdanove AJ, Wise RP: **Convergent evolution of effector protease recognition by *Arabidopsis* and barley.** *Mol Plant Microbe Interact* 2019, **32**:550–565, <https://doi.org/10.1094/MPMI-07-18-0202-FI>.
 36. Kohler A, Kuo A, Nagy LG, et al.: **Convergent losses of decay mechanisms and rapid turnover of symbiosis genes in mycorrhizal mutualists.** *Nat Genet* 2015, **47**:410–415, <https://doi.org/10.1038/ng.3223>.
 37. Delaux PM, Radhakrishnan GV, Jayaraman D, Cheema J, Malbreil M, Volkening JD, Sekimoto H, Nishiyama T, Melkonian M, Pokorny L, Rothfels CJ, Sederoff HW, Stevenson DW, Surek B, Zhang Y, Sussman MR, Dunand C, Morris RJ, Roux C, Wong GK, Oldroyd GE, Ané JM: **Algal ancestor of land plants was preadapted for symbiosis.** *Proc Natl Acad Sci U S A* 2015, **112**:13390–13395.
 38. Hong JJ, Park YS, Bravo A, et al.: **Diversity of morphology and function in arbuscular mycorrhizal symbioses in Brachypodium distachyon.** *Planta* 2012, **236**:851–865.
 39. Liu F, Xu Y, Jiang H, Jiang C, Du Y, Gong C, Wang W, Zhu S, Han G, Cheng B: **Systematic identification, evolution and expression analysis of the Zea mays PHT1 gene family reveals several new members involved in root colonization by arbuscular mycorrhizal fungi.** *Int J Mol Sci* 2016, **17**:930.
 40. Boch J, Scholze H, Schornack S, Landgraf A, Hahn S, Kay S, Lahaye T, Nickstadt A, Bonas U: **Breaking the code of DNA binding specificity of TAL-type III effectors.** *Science* 2009, **326**:1509–1512.
 41. Moscou MJ, Bogdanove AJ: **A simple cipher governs DNA recognition by TAL effectors.** *Science* 2009, **326**:1501.
 42. de Lange O, Wolf C, Dietze J, Elsaesser J, Morbitzer R, Lahaye T: **Programmable DNA-binding proteins from Burkholderia provide a fresh perspective on the TALE-like repeat domain.** *Nucleic Acids Res* 2014, **42**:7436–7449.
 43. Erkes A, et al.: **Evolution of transcription activator-like effectors in *Xanthomonas oryzae*.** *Genome Biol Evol* 2017, **9**:1599–1615.
 44. Doyle EL, Stoddard BL, Voytas DF, Bogdanove AJ: **TAL effectors: highly adaptable phyto-bacterial virulence factors and readily engineered DNA-targeting proteins.** *Trends Cell Biol* 2013, **23**:390–398.
 45. Hutin M, Pérez-Quintero AL, Lopez C, Szurek B: **MorTAL: Kombat: the story of defense against TAL effectors through loss-of-susceptibility.** *Front Plant Sci* 2015, **6**:535.
 46. Teper D, Wang N: **Consequences of adaptation of TAL effectors on host susceptibility to *Xanthomonas*.** *PLoS Genet* 2021, **17**, e1009310.
 47. Sarris PF, Cevik V, Dagdas G, Jones JD, Krasileva KV: **Comparative analysis of plant immune receptor architectures uncovers host proteins likely targeted by pathogens.** *BMC Biol* 2016, **14**:8.
 48. Shimizu M, Hirabuchi A, Sugihara Y, Abe A, Takeda T, Kobayashi M, Hiraka Y, Kanzaki E, Oikawa K, Saitoh H, Langner T, Banfield MJ, Kamoun S, Terauchi R: **A genetically linked pair of NLR immune receptors show contrasting patterns of evolution.** *Proc Natl Acad Sci U S A* 2022, **119**, e2116896119.
This study illustrates the incredibly dynamic evolution of integrated domains in NLR immune receptors and the contrasting patterns of diversification between paired NLRs.
 49. Maqbool A, Saitoh H, Franceschetti M, Stevenson CEM, Uemura A, Kanzaki H, Kamoun S, Terauchi R, Banfield MJ: **Structural basis of pathogen recognition by an integrated HMA domain in a plant NLR immune receptor.** *eLife* 2015, **4**, e08709.
 50. Bialas A, Zess EK, De la Concepcion JC, Franceschetti M, Pennington HG, Yoshida K, Upson JL, Chanclud E, Wu C-H, Langner T, Maqbool A, Varden FA, Derevnina L, Belhaj K, Fujisaki K, Saitoh H, Terauchi R, Banfield MJ, Kamoun S: **Lessons in effector and NLR biology of plant-microbe systems.** *Mol Plant Microbe Interact* 2018, **31**:34–45.

51. De La Concepcion JC, Benjumea JV, Bialas A, Terauchi R, Kamoun S, Banfield MJ: **Functional diversification gave rise to allelic specialization in a rice NLR immune receptor pair.** *eLife* 2021, **10**, e71662.
52. Kourelis J, Marchal C, Posbeyikian A, Harant A, Kamoun S: **NLR immune receptor-nanobody fusions confer plant disease resistance.** *Science* 2023, **379**:934–939.
53. Bentham AR, De la Concepcion JC, Vega Benjumea J, Jones S, Mendel M, Stubbs J, Stevenson CEM, Maidment JHR, Youles M, Kourelis J, Zdrzalek R, Kamoun S, Banfield MJ: **Allelic compatibility in plant immune receptors facilitates engineering of new effector recognition specificities.** *Plant Cell* 2023, koad204.
54. Arnold FH: **Innovation by evolution: bringing new chemistry to life.** *Angew Chem Int Ed* 2019, **58**:14420–14426.