

An ancestral oomycete locus contains late blight avirulence gene *Avr3a*, encoding a protein that is recognized in the host cytoplasm

Miles R. Armstrong*, Stephen C. Whisson*, Leighton Pritchard*, Jorunn I. B. Bos[†], Eduard Venter*, Anna O. Avrova*, Anne P. Rehmany[‡], Ulrike Böhme[§], Karen Brooks[§], Inna Cherevach[§], Nancy Hamlin[§], Brian White[§], Audrey Fraser[§], Angela Lord[§], Michael A. Quail[§], Carol Churcher[§], Neil Hall[§], Matthew Berriman[§], Sanwen Huang[¶], Sophien Kamoun[†], Jim L. Beynon[‡], and Paul R. J. Birch*^{||}

*Plant Pathogen Interactions Programme, Scottish Crop Research Institute, Invergowrie, DD2 5DA Dundee, United Kingdom; [†]Department of Plant Pathology, Ohio Agricultural Research and Development Center, Ohio State University, Wooster, OH 44691; [‡]Warwick HRI, University of Warwick, Wellesbourne, Warwick CV35 9EF, United Kingdom; [§]The Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SA, United Kingdom; and [¶]Laboratory of Plant Breeding, Department of Plant Sciences, Graduate School of Experimental Plant Sciences, Wageningen University, Binnenhaven 5, 6709 PD, Wageningen, The Netherlands

Edited by R. James Cook, Washington State University, Pullman, WA, and approved April 11, 2005 (received for review January 6, 2005)

The oomycete *Phytophthora infestans* causes late blight, the potato disease that precipitated the Irish famines in 1846 and 1847. It represents a reemerging threat to potato production and is one of >70 species that are arguably the most devastating pathogens of dicotyledonous plants. Nevertheless, little is known about the molecular bases of pathogenicity in these algae-like organisms or of avirulence molecules that are perceived by host defenses. Disease resistance alleles, products of which recognize corresponding avirulence molecules in the pathogen, have been introgressed into the cultivated potato from a wild species, *Solanum demissum*, and *R1* and *R3a* have been identified. We used association genetics to identify *Avr3a* and show that it encodes a protein that is recognized in the host cytoplasm, where it triggers *R3a*-dependent cell death. *Avr3a* resides in a region of the *P. infestans* genome that is colinear with the locus containing avirulence gene *ATR1^{NdwsB}* in *Hyaloperonospora parasitica*, an oomycete pathogen of *Arabidopsis*. Remarkably, distances between conserved genes in these avirulence loci were often similar, despite intervening genomic variation. We suggest that *Avr3a* has undergone gene duplication and that an allele evading recognition by *R3a* arose under positive selection.

microsynteny | *Phytophthora infestans* | hypersensitive response | linkage disequilibrium | hemibiotrophy

Phytophthora *infestans* causes late blight, the devastating potato disease responsible for the Irish famines in the mid-1840s (1). For decades, disease control has involved regular applications of agrochemicals, although recent widespread occurrence of new fungicide-resistant strains has led many to consider this pathogen as a reemerging threat to global food security (2, 3). It is one of >70 *Phytophthora* species that are arguably the most devastating pathogens of dicotyledonous plants. Other economically important phytophthoras include *Phytophthora sojae*, cause of soybean root rot; *Phytophthora palmivora*, cause of cocoa black pod; *Phytophthora cinnamomi*, cause of dieback and root rot in >2,000 plant species; and the recently identified *Phytophthora ramorum* that is decimating oak trees in the United States and throughout Europe (4). Despite a fungus-like filamentous growth habit, they are related to brown algae in the kingdom Stramenopiles (5). Whereas fungal pathogenicity has been intensively studied, little is known about the molecular genetics of oomycete pathogenicity or about pathogen molecules that are recognized by host defenses (1, 4).

Over the last 75 years, potato breeders have introduced at least 11 late blight resistance (*R*) alleles from *Solanum demissum* into the cultivated potato (6). The products of *R* alleles recognize the products of corresponding avirulence (*Avr*) alleles in races of *P.*

infestans, triggering disease resistance and a localized programmed cell death called the hypersensitive response (HR). Recently, the potato *R3* locus was shown to contain two tightly linked genes, *R3a* and *R3b*, with distinct specificities (7). *R3a* has since been cloned and encodes a presumed cytoplasmic coiled-coil nucleotide binding site leucine-rich repeat protein (8). Although several *P. infestans* *Avr* loci have been genetically mapped (1, 9), no *Avr* genes have yet been reported, mainly due to difficulties encountered with positional cloning, such as high levels of repetitive DNA and aberrant segregation at the target locus. Our knowledge of the molecular basis of race evolution in this pathogen is thus limited.

Many AVR proteins from fungal plant pathogens possess N-terminal type II signal peptides (SPs) for secretion and exhibit significant sequence variation between pathogen races. Thus, *Avr* alleles in virulent races of the tomato pathogen *Cladosporium fulvum* (10, 11) and the barley pathogen *Rhynchosporium secalis* (12) contain SNPs yielding secreted proteins that do not elicit HR on plants containing corresponding *R* alleles. We therefore predicted (13) that a candidate gene approach of screening ESTs for genes encoding secreted proteins, followed by association genetic [linkage disequilibrium (LD)] studies, would be an alternative route to identify *P. infestans* *Avr* genes.

A strategy for predicting *Phytophthora* extracellular protein (*pex*) genes was reported that identified 142 such genes from *P. infestans* (14). One EST, *Pex147* (encoding a 147-aa protein; GenBank accession no. BE776395), was similar to a *P. sojae* gene, *avr1b*, encoding an elicitor of HR in soybean plants containing *Rps1b* (15). The PEX147 protein product was found in *P. infestans* culture filtrate, demonstrating that it was secreted (14). We used an association genetics approach to show that *Pex147* was likely to be *Avr3a* in *P. infestans* and confirmed that this was the case by transient expression of alleles of the gene in potato genotypes containing a variety of *R* genes and by coexpression of *Avr3a* and *R3a* in *Nicotiana benthamiana*.

Materials and Methods

***P. infestans* Isolates and Analyses of Polymorphism.** *P. infestans* isolates were from the Scottish Crop Research Institute culture

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: SP, signal peptide; *Avr*, avirulence; LD, linkage disequilibrium; BAC, bacterial artificial chromosome; CDS, coding sequence; HR, hypersensitive response; *R*, resistance; *pex*, *Phytophthora* extracellular protein.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database [accession nos. AC146942 (Pi-BAC-61F2), AJ893356 (Pi-BAC-35J4), and AJ893357 (Pi-BAC-49P21)].

^{||}To whom correspondence should be addressed. E-mail: pbirch@sri.sari.ac.uk.

© 2005 by The National Academy of Sciences of the USA

collection (Table 1, which is published as supporting information on the PNAS web site). DNA was prepared by using the DNeasy plant minikit (Qiagen, Valencia, CA). PCR primers (Pex147F, 5'-CCATGCGTCTGGCAATTATGCT-3'; Pex147R, 5'-CTGAAAATAATATCCAGTGA-3') were used to amplify Pex147 from each isolate. PCR products were purified by using the QIAquick PCR purification kit (Qiagen) and were directly sequenced by using the Applied Biosystems BigDye v3.1 Terminator sequencing kit or used to generate a radiolabeled probe for hybridization to the *P. infestans* bacterial artificial chromosome (BAC) library (see below).

Analyses of *P. infestans* BAC Clones, BAC Sequencing, and Annotation. Hybridization of coding sequences (CDSs) (PCR-amplified by using primers shown in Table 2, which is published as supporting information on the PNAS web site) and Pex147 to the *P. infestans* BAC library and BAC DNA preparations were as described in ref. 16. Pi-BAC-61F2 (GenBank accession no. AC146942) was sequenced at the Broad Institute (Cambridge, MA). Pi-BAC-35J4 (accession no. AJ893356) and Pi-BAC-49P21 (accession no. AJ893357) were sequenced at The Wellcome Trust Sanger Institute. BAC sequencing and assembly were essentially performed as described in ref. 17. Clones were sequenced to 8-fold coverage and finished.

Annotation was performed by using ARTEMIS software (18). Functional assignments were based on assessment of BLAST and FASTA searches against public databases, and for domain predictions, searches such as INTERPRO (19), TMHMM 2.0 (20), and SIGNALP 2.0 (www.cbs.dtu.dk/services/SignalP) were carried out. The comparison of the avirulence loci of *Hyaloperonospora parasitica* and *P. infestans* was carried out by using BLAST comparisons viewed with the ARTEMIS comparison tool (K. Rutherford, unpublished).

Analyses of Transcript Levels. Maintenance of *P. infestans* isolate 88069; preparation of nonsporulating mycelium, sporangia, zoospores, germinating cysts, and germinating cysts developing appressoria; and inoculations of potato were carried out as described in refs. 21 and 22. For real-time RT-PCR, material was prepared, RNA extracted, and experiments conducted as described previously by using the *actA* gene from *P. infestans* as a constitutively expressed endogenous control (21). Real-time RT-PCR primers for Pex147 were 5'-CGCCATAAACTTTGCAACCA-3' and 5'-TGCCGGCTGAATCGTGTAT-3' (amplicon size = 92 bp), primers for Pex147-2 were 5'-GGTGGCAGCACAAGAGGC-3' and 5'-GCAGCTATTGTAGATCCGATTGTATC-3' (amplicon size = 104 bp), and primers for Pex147-3 were 5'-TTGGTGGCAGCACAATGGT-3' and 5'-CCCAGGTGCATCAGGTAGCT-3' (amplicon size = 120 bp). Repeated amplifications, on independent occasions with different cDNA samples, resulted in similar expression profiles.

Expression Constructs, Particle Cobombardment, and Agrobacterium-Infiltration Experiments. The *pex147* ORF was PCR-amplified from heterozygous *P. infestans* isolate SC95-1.1.2 (Table 1) by using primers 5'-CACACCATGGGTCTGGCAATTA-3' (introducing an NcoI site at the 5' end) and 5'-GTTTCAGCATCTAGAATCGGATTTTCTG-3' (introducing an XbaI site at the 3' end) and cloned into pRTL2 (23) for particle cobombardment and primers 5'-GGAAATCGATTCTCTCAGCTC-CCCAGGGTTCCAC-3' (introducing a ClaI site at the 5' end) and 5'-GGAAGCGGCCGCCACGAGCGTTTCAGCAGTTAGAATCGG-3' (introducing a NotI site at the 3' end) for cloning into potato virus X-*Agrobacterium tumefaciens* binary vector pGR106 (24). The alleles were cloned without SP-encoding sequences into pRTL2 by using additional primer 5'-GGAACCATGGACCAAACCAAGGTCTGG-3' and into pGR106 by using additional primer 5'-GGAAATCGATATG-

GACCAAACCAAGGTCTGGT-3', introducing methionine in place of isoleucine at position 22 [predicted to be the SP cleavage site by using SIGNALP (www.cbs.dtu.dk/services/SignalP)]. PCR products digested with NcoI and XbaI or ClaI and NotI were ligated into pRTL2 or pGR106 (respectively), electroporated into Electromax DH10B *Escherichia coli*, and selected on LB ampicillin (pRTL2) or LB kanamycin/tetracycline (pGR106) by using standard protocols. Individual clones were sequenced to ensure that errors had not been introduced.

Biolistic cobombardment was performed by using the Bio-Rad PDS-1000/He Biolistic Particle Delivery System. One-micrometer gold particles were coated with 0.83 $\mu\text{g}/\text{mg}$ each pRTL2::*gfp* and pRTL2 expressing test sequences. Leaves from 4- to 6-week-old potato plants were cut down the midrib. Half-leaves from cv. Bintje or cv. Craigs Royal lacking either *R* gene-mediated (*r*) or field resistances to *P. infestans* were cobombarded alongside half-leaves from potato genotypes containing *R3a* (cv. Pentland Ace), *R3b* (SW8563-0182), *R1* (cv. Craigs Snow White), *R2* (1512c[16]), or *R10* (3681ad[1]) to receive similar quantities of particles. Transgenic cv. Desireé containing *R3a* (T68.4-002) was similarly screened with adjacent nontransgenic cv. Desireé, which, again, lacks either *R* gene-mediated or field resistances to *P. infestans*. Half-leaf pairs were cobombarded at 1,550 psi and incubated in light for 20–24 h at 20°C followed by 18–24 h in dark at 20°C. The half-leaves were imaged by using a Bio-Rad MRC 1000 confocal laser scanning microscope, and GFP was excited by using blue laser light at 488 nm. Numbers of GFP-fluorescing cells were quantified. For each half-leaf pair, the percentage of GFP fluorescence was determined, and the figure for the *R* gene-containing cultivar was divided by that observed on Bintje, yielding a GFP fluorescence ratio. For each line, three leaf pairs from independent plants were cobombarded with pRTL2::*gfp* and pRTL2 expressing each test sequence. A further three leaf pairs were bombarded with particles coated with pRTL2::*gfp* alone. These experiments were repeated, and the means and standard deviations are presented in Fig. 4.

Agrobacterium infiltration was performed in *N. benthamiana* by using *Agrobacterium tumefaciens* strain AGL0 carrying the pBINplus::*R3a* construct (8) and strain LBA4404 carrying the pGR106::*Avr3a*-K⁸⁰ I¹⁰³ and pGR106::*Avr3a*-E⁸⁰ M¹⁰³ (both without SP) constructs. Recombinant *A. tumefaciens* cultures were grown and induced before infiltration as described in ref. 25, except that culturing was performed in LB supplemented with 50 μM kanamycin. *A. tumefaciens* cultures carrying the *R3a* and *Avr3a* constructs were mixed in a 2:1 ratio before centrifugation. For transient coexpression, the harvested cells were resuspended in MMA buffer containing 200 μM acetosyringone (25) to a final OD of 0.4. [One liter of MMA contains 5 g of MS salts (Duchefa, Haarlem, The Netherlands), 1.9 g of MES (*N*-morpholinoethanesulfonic acid; Sigma), and 20 g of sucrose, pH adjusted to 5.6 with 1 M NaOH (25).] For transient expression of the *Avr3a* genes, cells harvested from *A. tumefaciens* cultures carrying the *Avr3a*-K⁸⁰ I¹⁰³ or *Avr3a*-E⁸⁰ M¹⁰³ constructs were resuspended to a final OD of 0.133 and infiltrated into 5- to 6-week-old leaves with a 3-ml syringe. Symptom development was monitored starting from 3 days after infiltration.

Results and Discussion

Identification of *Avr3a* Through Association Genetics. Primers were designed to PCR-amplify Pex147 from 55 *P. infestans* strains isolated from countries in Europe and North and South America that had been virulence-tested on potato genotypes containing 9 of the 11 *S. demissum* *R* genes. After sequencing the PCR products, only three SNPs were found, changing amino acids S19C, E80K, and M103I (Fig. 1). These SNPs revealed only two alleles that showed 100% correlation with virulence phenotype




Fig. 1. Protein alignments and phylogeny of AVR3a, avr3a, and PEX147-like paralogues. (A) Multiple alignment (using www.ebi.ac.uk/clustalw) of predicted proteins (AVR3a and avr3a) derived from the *Avr3a* (avirulent) and *avr3a* (virulent) alleles (initially termed *Pex147*) with those from the two paralogous *P. infestans* *Pex147-2* and *Pex147-3* sequences and the AVR1b protein from *P. sojae*. The N-terminal type II SP (first 21 aa) and the RXLR motif, found also in the *H. parasitica* ATR1^{NdVsb} protein (32), are underlined. The locations of the three amino acid polymorphisms are indicated by numbers above the amino acids. (B) Maximum likelihood estimation of the phylogeny of *Avr3a*, *avr3a*, and *Pex147*-like paralogues using a nucleotide alignment based on the amino acid alignment in A. The tree was constructed by using DNAML from the PHYLIP package, with the *P. sojae* *Avr1b* sequence (AAM20934) as an outgroup. Bootstrap values from 1,000 random resamplings of the data are indicated. The alignment supports the closer evolutionary relationship of the *Avr3a* allele, rather than the *avr3a* allele, to the flanking *Pex147-2* and *Pex147-3* paralogues.

on cv. Pentland Ace, containing *R3a* (Table 1). Presence of the C¹⁹ K⁸⁰ I¹⁰³ (C-K-I) allele was associated with avirulence, whereas virulent isolates were homozygous for the S¹⁹ E⁸⁰ M¹⁰³ (S-E-M) allele.

The possibility that the observed association resulted from genetic hitchhiking (26) was considered; *Pex147* could simply be closely linked to *Avr3a* rather than the gene itself. In sexual populations, hitchhiking decreases with physical distance because of recombination. After hybridization of *Pex147* to a *P. infestans* BAC library (16), BAC clones were identified from which SNPs flanking the gene were sought to investigate the region of LD. The sequence of Pi-BAC-49P21 revealed *Pex147* and two additional *Pex147*-like sequences (Fig. 1). A further *Pex147*-like sequence contained a frameshift mutation and was annotated as a pseudogene (Fig. 2). The occurrence of SNPs 9.6 kb upstream and 10.9 kb downstream of *Pex147* (Fig. 2) in 10 *P. infestans* isolates showed no association with the *Avr3a*-*R3a* phenotype (Table 1), indicating breakdown in LD and ruling out the *Pex147-2* and *Pex147-3* paralogues as candidate *Avr3a* genes. Within the 20-kb region defined by these SNPs, only four additional CDSs were identified (results not shown), none of which encoded proteins with SPs, and *Pex147* was thus selected as the most likely *Avr3a* candidate.

***Pex147* Is Up-Regulated Before and During Potato Infection.** Real-time RT-PCR was used to investigate the expression of *Pex147* and the flanking *Pex147-2* and *Pex147-3* paralogues in *P. infestans* preinfection cell types (sporangia, zoospores, germinating cysts, and germinating cysts forming appressoria) and infected susceptible potato cv. Bintje 12, 24, 33, 48, 56, and 72 h postinoculation (hpi). Expression of each gene was compared with the *ActA* gene from *P. infestans*, which has been used as a constitutively expressed endogenous control (21, 22). Although PCR products of expected sizes were generated for *Pex147-2* and *Pex147-3* from genomic DNA template, no products were detected from cDNA templates in the case of *Pex147-3*, and expression was only weakly detected in the case of *Pex147-2* and

only in cDNA from axenic cultured, nonsporulating mycelium. In contrast, expression of *Pex147* was readily detectable in preinfection and infection stages and, in these samples, was compared with the level of its expression in a calibrator sample, cDNA from cultured nonsporulating mycelium, which was assigned the value of 1.0. *Pex147* was up-regulated >100-fold in germinating cysts developing appressoria and showed elevated levels of expression throughout infection, with an early peak of expression at 24 hpi, in the biotrophic phase of infection, and >200-fold elevation of expression at 72 hpi, in the necrotrophic phase of the interaction (Fig. 3). Repeated amplifications on independent occasions with different cDNA samples resulted in similar expression profiles. The up-regulation of *Pex147* immediately before and during potato infection supports a potential role in pathogenicity. In contrast, the failure to detect such expression of the flanking *Pex147-2* and *Pex147-3* paralogues supports the LD analysis, ruling them out as *Avr3a* candidates.

Coexpression in Potato of *Pex147* Alleles with *gfp* Demonstrates That *Pex147* Is *Avr3a*. The S-E-M (virulent) and C-K-I (avirulent) alleles of *Pex147* and each allele truncated to remove SP-encoding sequences (and thus lacking the S19C polymorphism) were transiently expressed in potato genotypes lacking a late blight *R* gene (cv. Bintje or cv. Craigs Royal) or containing *R3a*, *R3b*, *R1*, *R2*, and *R10*. In each case, transient expression of the alleles involved biolistic bombardment with a second vector expressing *gfp* as a marker of cell vitality. Similar approaches were used to indicate and quantify HR triggered by *Avr* genes from the bacterium *Pseudomonas syringae* (27), the fungus *Magnaporthe grisea* (28), and the oomycete *H. parasitica* (29). There was no reduction in GFP fluorescence on leaves from any plants cobombarded with alleles possessing SP-encoding sequences, relative to that observed when GFP was bombarded alone (data not shown). Furthermore, no difference in GFP fluorescence was seen in leaves from Bintje or Craigs Royal compared with plants containing *R3b*, *R1*, *R2*, or *R10* that were cobombarded with alleles lacking SP-encoding sequences (Fig.




Fig. 4. Cobombardment demonstrates recognition of AVR3a by plants containing R3a. Truncated ($E^{80} M^{103}$) *avr3a* and ($K^{80} I^{103}$) *Avr3a* alleles were transiently coexpressed with *gfp* in potato genotypes containing *R1* (a), *R2* (b), *R3b* (c), *R10* (d), and *R3a* (e), each compared with cv. Bintje (f), and in transgenic cv. Desiree expressing *R3a* compared with cv. Desiree (f). Typically, more GFP-expressing cells were observed on *R* gene-containing genotypes when *gfp* was bombarded alone than on adjacent Bintje leaves (a–e), resulting in GFP fluorescence indexes > 1 (where 1 is the average total GFP fluorescence on Bintje). In each histogram, the average GFP fluorescence index is presented after coexpression with *Avr3a* (light gray) or *avr3a* (dark gray) and expression of *gfp* alone (white).

tration (15). However, it is possible that infiltration of large quantities of protein could lead to uptake of AVR1b into the host cell through endocytosis.

Analysis of *Avr3a* and *ATR1^{NdWsB}* Loci Reveals Conservation of Microsynteny. Analysis of >250 kb spanning *Avr3a* revealed a surprising finding: Several CDSs were strikingly similar to CDSs flanking *ATR1^{NdWsB}* (32) in *H. parasitica* (Table 2), suggesting similar gene content in these avirulence loci. Genes encoding methylenetetrahydrofolate dehydrogenase (MTD), a RAS-like protein, F-actin capping protein, and 3-isopropylmalate dehydratase (3-IPMDH) (Fig. 2) were colinear in the two loci and

similar distances apart, although they were apparently inverted relative to other genes. It is interesting to note that MTD- and dimeric dihydrodiol dehydrogenase-encoding sequences are duplicated in *P. infestans*, and this duplication may have coincided with inversion (Fig. 2). Three intact copies of *Pex147* and a *Pex147*-like pseudogene again indicate gene duplication in *P. infestans*, and, if extended with similar frequency throughout the genome, such duplication would contribute significantly to global gene redundancy. This phenomenon is marked in the yeast *Debaryomyces hansenii* and distinguishes it from other yeasts (33). Three Lon protease (LON)-encoding sequences in *H. parasitica*, two of which are partial sequences and, therefore, possibly pseudogenes, indicate that gene duplication is not exclusive to *P. infestans* in this ancestral locus (Fig. 2).

ATR1^{NdWsB} (32) and *Avr3a* share little sequence similarity but are in strikingly similar locations in their respective loci and are flanked on one side by one or more LON-encoding sequences and on the other side by vacuolar proton ATPase subunit (H+ PUMP)- and myb-encoding genes (Fig. 2). The H+ PUMP- and myb-encoding genes are again similar distances apart and are also transcriptionally inverted, suggesting numerous structural rearrangements in the *Avr3a* and *ATR1^{NdWsB}* loci. Remarkably, the distances between intact LON sequences and the myb- or H+ PUMP-encoding sequences are also similar, suggesting that an unknown mechanism of positional conservation has acted at sites within these oomycete genomes, between which are regions of variation. A similar phenomenon has been noted in studies of conservation of synteny between rice and *Arabidopsis thaliana* (34). Although recently grouped together in the Peronosporales based on molecular phylogenies (35), *P. infestans* and *H. parasitica* are not particularly closely related (36). Considering the high sequence diversity observed in the Peronosporales, the colinearity is therefore notable.

Within these colinear loci, the avirulence genes have apparently evolved differently. In *H. parasitica*, considerable allelic variation was observed in *ATR1^{NdWsB}* (32). In *P. infestans*, gene duplication and divergence has provided a mechanism for generating variation in *Avr3a*-like sequences, and numerous synonymous and nonsynonymous nucleotide polymorphisms distinguish these genes. The *avr3a* allele is distinguished from the *Pex147-2* gene by 10 synonymous and 22 nonsynonymous polymorphisms and is distinguished from *Pex147-3* by 8 synonymous and 20 nonsynonymous polymorphisms. The *Pex147-2* and *Pex147-3* paralogues are distinguished from each other by 9 synonymous and 21 nonsynonymous polymorphisms (Fig. 1).




Fig. 5. Transient expression and coexpression of *Avr3a* and *R3a* in *N. benthamiana*. Leaves of *N. benthamiana* plants were infiltrated with *A. tumefaciens* carrying pGR106::*Avr3a*-K-I or pGR106::*avr3a*-E-M (expressing mature forms of the proteins lacking SPs) alone or mixed with an *A. tumefaciens* strain carrying pBINplus::*R3a*. For transient coexpression of *Avr3a* and *R3a*, the *A. tumefaciens* solutions were mixed in a 1:2 ratio before infiltration. Photographs of symptoms were taken 6 days postinfiltration. Circles indicate the infiltrated area on the leaf panels for each treatment. This is a representative leaf from multiple assays and experiments.

Nevertheless, only two alleles of *Avr3a* were observed in 55 *P. infestans* individuals, and the SNPs distinguishing them all result in nonsynonymous amino acid substitutions, suggesting that diversifying selection, perhaps driven by coevolution with host plants, has acted on this gene. Interestingly, diversifying selection has also acted on the *R3a* gene (8) and its paralogues, suggesting a coevolutionary “arms race” between *R3a* and *Avr3a*. The amino acids specific to AVR3a (C¹⁹ K⁸⁰ I¹⁰³) are shared with PEX147-2, and two are shared with PEX147-3. The absence of other SNPs, or of selectively neutral SNPs between *Avr3a* and *avr3a*, suggests that the virulent allele arose from an avirulent progenitor sequence under positive selection (Fig. 1).

The allelic diversity in the *ATRI*^{NdWsb} (32) and *Avr3a* genes are in stark contrast, and it should be noted that *H. parasitica* isolates were derived from natural populations coevolving with wild *A. thaliana*, whereas *P. infestans* isolates were largely obtained from cultivated potato in countries distant from Mexico, a center of diversity for the pathogen. It is well documented that *P. infestans* underwent a series of extreme genetic bottlenecks during its pan-global distribution (37), and a lack of allelic diversity in *Avr3a* may reflect this. It will thus be interesting to compare *Avr3a* allele frequency and diversity in natural Mexican *P. infestans* populations with those found in agricultural populations elsewhere.

Conclusions

We report the use of association genetics to identify the *P. infestans Avr3a* gene and show that its product is recognized in

an *R3a*-dependent manner in the host cytoplasm. Analysis of the *Avr3a* locus revealed unexpected conservation of synteny with the locus containing *ATRI*^{NdWsb} in *H. parasitica*. This colinearity supports comparative genomics as an approach to investigate the evolution of pathogenicity in oomycetes. The isolation of *R3a* (8) and *Avr3a* represents an opportunity for detailed investigation of the earliest recognition events in a potato-*P. infestans* R-AVR interaction and of subsequent signaling pathways leading to disease resistance. It also opens a door to studies of molecular mechanisms potentially underlying the biotrophic and necrotrophic phases of the *P. infestans* infection cycle. The current economic importance of this pathogen and its roles in the history of Ireland and in the establishment of the field of plant pathology means that this investigation is a significant step forward in applied and fundamental late blight research.

We thank Vivianne Vleeshouwers and Richard Visser (both of Wageningen University) for tissue samples (T68.4-002 and SW8563-0182) and Petra Boevink (Scottish Crop Research Institute) for pRTL2::gfp and for critically reading the manuscript. This work was supported by the Scottish Executive Environment and Rural Affairs Department, the Biotechnology and Biological Sciences Research Council, and the National Science Foundation. J.I.B.B. and S.K. were supported by National Science Foundation Plant Genome Research Program Grant DBI-0211659.

- Birch, P. R. J. & Whisson, S. C. (2001) *Mol. Plant Pathol.* **2**, 257–263.
- Schiermeier, Q. (2001) *Nature* **410**, 1011.
- Duncan, J. M. (1999) *Microbiol. Today* **26**, 114–116.
- Kamoun, S. (2003) *Eukaryot. Cell* **2**, 191–199.
- Sogin, M. L. & Silberman, J. D. (1998) *Int. J. Parasitol.* **28**, 11–20.
- Gebhardt, C. & Valkonen, J. P. T. (2001) *Annu. Rev. Phytopathol.* **39**, 79–102.
- Huang, S., Vleeshouwers, V. G. A. A., Werij, J. S., Hutten, R. C. B., van Eck, H. J., Visser, R. G. F. & Jacobsen, E. (2004) *Mol. Plant–Microbe Interact.* **17**, 428–435.
- Huang, S., van der Vossen, E. A. G., Huang, H., Vleeshouwers, V. G. A. A., Zhang, N., Borm, T. J. A., van Eck, H. J., Baker, B., Jacobsen, E. & Visser, R. (2005) *Plant J.* **42**, 251–261.
- van der Lee, T., Robold, A., Testa, A., van 't Klooster, J. W. & Govers, F. (2001) *Genetics* **157**, 949–956.
- Joosten, M. H. A. J., Vogelsang, R., Cozignsen, T. J., Verberne, M. C. & de Wit, P. J. G. M. (1997) *Plant Cell* **9**, 367–379.
- Luderer, R., Takken, F. L., de Wit, P. J. G. M. & Joosten, M. H. (2002) *Mol. Microbiol.* **45**, 875–884.
- Rohe, M., Gierlich, A., Hermann, H., Hahn, M., Schmidt, B., Rosahl, S. & Knogge, W. (1995) *EMBO J.* **14**, 4167–4177.
- Bos, J. I. B., Armstrong, M., Whisson, S. C., Torto, T. A., Ochwo, M., Birch, P. R. J. & Kamoun, S. (2003) *New Phytol.* **159**, 63–72.
- Torto, T. A., Li, S., Styer, A., Huitema, E., Testa, A., Gow, N. A. R., van West, P. & Kamoun, S. (2003) *Genome Res.* **13**, 1675–1685.
- Shan, W., Cao, M., Leung, D. & Tyler, B. (2004) *Mol. Plant–Microbe Interact.* **17**, 394–403.
- Whisson, S. C., van der Lee, T., Bryan, G., Waugh, R., Govers, F. & Birch, P. R. J. (2001) *Mol. Genet. Genomics* **266**, 289–295.
- Harris, D. E. & Murphy, L. (2001) in *Methods in Molecular Biology*, eds Starkey, M. P. & Elaszwarapu, R. (Humana, Totowa, NJ), pp. 217–234.
- Rutherford, K., Parkhill, J., Crook, J., Horsnell, T., Rice, P., Rajandream, M. A. & Barrell, B. (2000) *Bioinformatics* **16**, 944–945.
- Apweiler, R., Attwood, T. K., Bairoch, A., Bateman, A., Birney, E., Biswas, M., Bucher, P., Cerutti, L., Corpet, F., Croning, M. D., et al. (2000) *Bioinformatics* **16**, 1145–1150.
- Krogh, A., Larsson, B., von Heijne, G. & Sonnhammer, E. L. (2001) *J. Mol. Biol.* **305**, 567–580.
- Avrova, A. O., Venter, E., Birch, P. R. J. & Whisson, S. C. (2003) *Fungal Genet. Biol.* **40**, 4–14.
- Liu, Z., Bos, J. I., Armstrong, M., Whisson, S. C., da Cunha, L., Torto-Alalibo, T., Win, J., Avrova, A. O., Wright, F., Birch, P. R. J. & Kamoun, S. (2005) *Mol. Biol. Evol.* **22**, 659–672.
- Restrepo, M. A., Freed, D. D. & Carrington, J. C. (1990) *Plant Cell* **2**, 987–998.
- Jones, L., Hamilton, A. J., Voinnet, O., Thomas, C. L., Maule, A. J. & Baulcombe, D. C. (1999) *Plant Cell* **11**, 2291–2301.
- Van der Hoorn, R. A., Laurent, F., Roth, R. & De Wit, P. J. (2000) *Mol. Plant–Microbe Interact.* **13**, 439–446.
- Barton, N. H. (2000) *Philos. Trans. R. Soc. London B* **355**, 1553–1562.
- Leister, R. T., Ausubel, F. M. & Katagiri, F. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 15497–15502.
- Jia, J., McAdams, S. A., Bryan, G. T., Hershey, H. P. & Valent, B. (2000) *EMBO J.* **19**, 4004–4014.
- Allen, R. L., Bittner-Eddy, P. D., Grenville-Briggs, L. J., Meitz, J. C., Rehmany, A. P., Rose, L. E. & Beynon, J. L. (2004) *Science* **306**, 1957–1960.
- Dodds, P. N., Lawrence, G. J., Catanzariti, A., Aycliffe, M. A. & Ellis, J. G. (2004) *Plant Cell* **16**, 755–768.
- Collmer, A., Lindeberg, M., Petnicki-Ocwieja, T., Schneider, D. J. & Alfano, J. R. (2002) *Trends Microbiol.* **10**, 462–469.
- Rehmany, A. P., Gordon, A., Allen, R. L., Armstrong, M. R., Whisson, S. C., Kamoun, S., Tyler, B. M., Birch, P. R. J. & Beynon, J. L. (2005) *Plant Cell*, in press.
- Dujon, B., Sherman, D., Fischer, G., Durrens, P., Caserogola, S., Lafontaine, I., de Montigny, J. Marck, C., Neuvéglise, C., Talla, E., et al. (2004) *Nature* **430**, 35–44.
- Han, F., Kilian, A., Chen, J. P., Kudrna, D., Steffenson, B., Yamamoto, K., Matsumoto, T., Sasaki, T. & Kleinjohs, A. (1999) *Genome* **42**, 1071–1076.
- Riethmüller, A., Voglmayr, H., Göker, M., Weiss, M. & Oberwinkler, F. (2002) *Mycologia* **94**, 834–849.
- Göker, M., Voglmayr, H., Riethmüller, A., Weiss, M. & Oberwinkler, F. (2003) *Can. J. Bot.* **81**, 672–683.
- Goodwin, S. B., Cohen, B. A. & Fry, W. (1994) *Proc. Natl. Acad. Sci. (USA)* **91**, 11591–11595.