



Review

Cite this article: Derevnina L *et al.* 2016

Emerging oomycete threats to plants and animals. *Phil. Trans. R. Soc. B* **371**: 20150459.

<http://dx.doi.org/10.1098/rstb.2015.0459>

Accepted: 15 August 2016

One contribution of 18 to a discussion meeting issue 'Tackling emerging fungal threats to animal health, food security and ecosystem resilience'.

Subject Areas:

plant science, microbiology

Keywords:

parasites, pathogens, disease, food security, agriculture, environment

Author for correspondence:

Sophien Kamoun

e-mail: sophien.kamoun@tsl.ac.uk

Emerging oomycete threats to plants and animals

Lida Derevnina¹, Benjamin Petre¹, Ronny Kellner², Yasin F. Dagdas¹, Mohammad Nasif Sarowar³, Artemis Giannakopoulou¹, Juan Carlos De la Concepcion¹, Angela Chaparro-Garcia¹, Helen G. Pennington¹, Pieter van West⁴ and Sophien Kamoun¹

¹The Sainsbury Laboratory, Norwich Research Park, Norwich NR4 7UH, UK

²Max Planck Institute for Plant Breeding Research, Carl-von-Linné-Weg 10, Cologne 50829, Germany

³Department of Fisheries Biology and Genetics, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

⁴International Centre for Aquaculture Research and Development, Aberdeen Oomycete Laboratory, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK

id LD, 0000-0001-7983-4187; BP, 0000-0002-3204-4249; RK, 0000-0002-4618-0110; YFD, 0000-0002-9502-355X; MNS, 0000-0001-7122-5299; AG, 0000-0002-7629-3108; JcDc, 0000-0002-7642-8375; HGP, 0000-0002-2980-9079; PwW, 0000-0002-0767-6017; SK, 0000-0002-0290-0315

Oomycetes, or water moulds, are fungal-like organisms phylogenetically related to algae. They cause devastating diseases in both plants and animals. Here, we describe seven oomycete species that are emerging or re-emerging threats to agriculture, horticulture, aquaculture and natural ecosystems. They include the plant pathogens *Phytophthora infestans*, *Phytophthora palmivora*, *Phytophthora ramorum*, *Plasmopara obducens*, and the animal pathogens *Aphanomyces invadans*, *Saprolegnia parasitica* and *Haliotidica noduliformans*. For each species, we describe its pathology, importance and impact, discuss why it is an emerging threat and briefly review current research activities.

This article is part of the themed issue 'Tackling emerging fungal threats to animal health, food security and ecosystem resilience'.

1. Introduction

Oomycetes, or water moulds, are fungal-like eukaryotes classified as stramenopiles, and are phylogenetically grouped with diatoms and brown algae [1–4]. They are among the most problematic group of disease-causing organisms in both agriculture and aquaculture, and represent a recurrent threat for global food security. Oomycetes cause some of the most devastating plant diseases affecting crops, ornamental plants and trees. They result in major economic losses and serious damage to natural ecosystems [5,6]. The most notorious species are members of the genus *Phytophthora*, such as the late blight pathogen *Phytophthora infestans*, known for triggering the Irish potato famine [7,8]. Other notable species include *Phytophthora palmivora* [9], causing cocoa black pod, and the sudden oak death pathogen *Phytophthora ramorum*, which emerged in more recent epidemics threatening native tree species [10]. Additional important oomycete plant pathogens include members of the *Pythium* genus and downy mildews [11,12].

In contrast with their terrestrial counterparts, aquatic oomycetes remain understudied [3,4]. This applies particularly to animal pathogenic oomycetes, such as *Saprolegnia* and *Aphanomyces* spp. [13]. Aquaculture has become one of the world's fastest growing food sectors, where freshwater fish dominate global aquaculture production [14]. Fish farming and fisheries provide livelihood and income for an estimated 54.8 million people, with employment in these sectors growing rapidly [15]. Diseases such as epizootic ulcerative syndrome and saprolegniosis, caused by oomycetes, are a significant threat to the aquaculture industry [16].

2. Why are oomycetes so successful?

The key to pathogenic oomycetes' success resides in their capacity to adapt to overcome host resistance and occasionally jump to new hosts. Various features of oomycete biology can explain their high evolutionary potential. One major driving force for their success is encompassed by their flexible mating system. Oomycetes can reproduce sexually (either homothallic or heterothallic), asexually or through interspecific hybridization [17] enabling them to gain both allelic diversity (sexual and interspecific hybridization) and to rapidly proliferate, resulting in large population sizes (asexual/clonal populations) [18]. Large population sizes compensates for the lack of sexual recombination and increase the occurrence of spontaneous mutants with enhanced fitness [18].

Genome sequencing of oomycetes has provided additional insights into their capacity to adapt to hosts. Oomycete genomes harbour large repertoires of genes encoding virulence effectors, which modulate host processes [19,20]. These effector genes are typically associated with fast evolving regions of the genome, particularly in areas enriched in repeats and transposable elements, promoting genome duplication, shuffling, increased rates of mutagenesis and gene silencing [21]. This bipartite genome organization, differentiated by slow and fast evolving regions, has led to the 'two-speed' genome model, where gene-sparse repeat-rich compartments serve as a cradle for adaptive evolution, underpinning infection success by enabling the pathogen to rapidly overcome host resistance, evolve new virulence determinants and even jump to new host species [22]. Another interesting aspect about oomycete genomes is that they exhibit variations in ploidy and can exist as either triploids or polyploids [23–25]. The relevance of this feature is currently not fully understood, however, polyploidy has been shown to enhance vigour and buffer mutational changes by masking deleterious alleles [26], and is therefore believed to play a vital role in pathogen success. In fact, many successful *P. infestans* clonal lineages are made up of triploid genotypes [26].

Here, we discuss seven oomycetes that are emerging or re-emerging threats to world agriculture, horticulture, aquaculture and natural ecosystems. They were selected based on their impact on economy and society, and include both well-known and understudied pathogens of both plants and animals. For each pathogen, we describe its pathology, importance and impact, discuss why it is an emerging threat and briefly review research activities that underpin effective disease management strategies. Table 1 summarizes our current knowledge of the genome and secreted proteins of the examined species and highlights why they are (re-)emerging threats.

3. *Phytophthora infestans*: the Irish potato famine pathogen

Phytophthora infestans infects potato and tomato plants, causing late blight disease. It is infamous for triggering the Irish potato famine in the mid-nineteenth century. Today, it is still a significant threat to global food security, causing severe yield losses in these crops, and economic losses of more than USD 6 billion annually [30,31]. In addition to destroying the foliage, *P. infestans* can infect tubers, which facilitates migration of the pathogen via the global seed tuber trade [32]. Current disease management strategies include chemical control and/or deployment of resistant cultivars. Resistance to agrochemicals,

however, is common in epidemic-causing isolates [33], and due to the pathogen's adaptability, disease resistance genes are also quickly broken down and rendered ineffective [34].

Disease outbreaks are typically caused by asexually reproducing clonal lineages [24,32,35,36]. These aggressive lineages dramatically reshape the population structure and become dominant over a short period of time [36,37]. For example, in Great Britain, genotype *13_A2*, first detected in July 2005, surged to more than 75% of the *P. infestans* population by 2008 (figure 1) [35]. Subsequently, *13_A2* was displaced by another genotype, *6_A1*, which increased in occurrence from 20% in 2010 to 80% in 2011 (<http://www.fwi.co.uk/academy/lesson/potatoes-understanding-blight1>). In addition, some *P. infestans* clonal lineages have become pandemic. *13_A2* was detected for the first time in South India in 2008, where it caused severe late blight epidemics in tomato, often resulting in 100% crop loss [38]. The emergence of this genotype in the Indian subcontinent has led to an increase in late blight incidence on tomato and potato, with epidemics reported in subsequent years following its introduction.

The *P. infestans* genome was sequenced in 2009 [8], which established this species as a model system, pioneering studies of oomycete effectors and evolution [8,40]. Notably, the study of *P. infestans* effectors have significantly moved forward our understanding of host processes targeted by the pathogen, forming strong foundations for plant–microbe interaction studies [19,41]. Furthermore, effectors have emerged as powerful tools to rapidly identify new resistances in potato germplasms.

4. *Phytophthora palmivora*: the pathogen of tropical plants

Phytophthora palmivora is a plant pathogen that infects more than 200 plant species in the tropics, including dicots and monocots [7]. Economically important hosts include cocoa, coconut, pineapple, rubber tree, durian, citrus, papaya and oil palm. Disease outbreaks in these crops impact the livelihood and nutrition of millions of people globally. Annual losses due to diseases caused by *P. palmivora*, such as oil palm bud rot and cocoa black pod, range from USD 250 million to over USD 1 billion, respectively [9].

Rainy season and high humidity favour pathogen sporulation leading to rapid disease progression and production of oospores. Disease spread from the soil into the canopy is often initiated through rain splash. Once the roots, leaves and fruits in the lower canopy are infected, the pathogen quickly spreads to the upper canopy by insects, wind and through wind-driven aerosols [7,42]. Additionally, transport of infected nursery plants, improper disposal of infected material, irrigation with zoospore-contaminated water, changing temperatures and global trade all contribute to the spread of *P. palmivora*. This makes it a serious threat to the economy of many developing countries, where cultivation of highly productive perennial crops is intensifying.

The broad host range of *P. palmivora* has enabled research using model plants such as *Medicago truncatula* [43], *Hordeum vulgare* (barley) [44] and *Nicotiana benthamiana* [45], moving forward the characterization of the molecular mechanisms involved in *P. palmivora* colonization. Notably, the process of root colonization by this pathogen is used to study similarities and differences between pathogenicity

Table 1. Features of oomycete pathogens highlighted in this paper, including the primary host, genome size, repetitive DNA content (as a percentage of genome size), number of protein coding genes, percentage of predicted secreted proteins and a summary of why the pathogen is a (re-)emerging threat. n.d., no data.

pathogen	important hosts (P, plant; A, animal)	common disease name (host)	why is it a (re-)emerging threat?	genome size (Mb) (% repetitive)	protein coding genes (% secreted) ^a	references
<i>Phytophthora infestans</i>	P—potato, tomato	late blight	new aggressive genotypes emerged in the UK (2008 and 2011) and India (2008)	240 (7.4)	18 155 (8.7)	[8]
<i>Phytophthora palmivora</i>	P—cocoa, coconut, pineapple, rubber tree, papaya, oil palm	bud rot (oil palm), black pod (cocoa), fruit rot (papaya)	outbreaks regularly destroy oil palm and cocoa plantations worldwide	n.d. (n.d.)	n.d. (n.d.)	—
<i>Phytophthora ramorum</i>	P—oak, tanoaks, rhododendron, beech, larch	sudden oak death (oak)	outbreaks destroyed forests in North America and Europe during the last 20 years	65 (17)	14 451 (10.5)	[27]
<i>Plasmopara obducens</i>	P— <i>Impatiens</i> species	<i>Impatiens</i> downy mildew	outbreaks occurred in nurseries worldwide within the last decade	70 (n.d.)	n.d. (n.d.)	[28]
<i>Aphanomyces invadans</i>	A—carp, perch, tilapia, snakeheads, salmonids, estuarine	epizootic ulcerative syndrome	regularly kills farmed and wild fish worldwide, threatening food safety in Asia and Africa	71 (n.d.)	15 248 (n.d.)	^b
<i>Saprolegnia parasitica</i>	A—fish (salmon), amphibians, crustaceans, aquatic insects	saprolegniosis	threatens fish farming in Europe, America and Asia, as well as endangered amphibian species worldwide	63 (17)	17 065 (5.7)	[29]
<i>Haliotida noduliformans</i>	A—abalone, mantis shrimp	abalone tubercle mycosis (abalone)	severe outbreaks occurred in South Africa and Japan between 2004 and 2006	n.d. (n.d.)	n.d. (n.d.)	—

^aIncluding putative effectors.^bSee http://www.ebi.ac.uk/ena/data/view/GCA_000520115.




Figure 1. An emerging clonal lineage reshaped *P. infestans* populations in Great Britain. (a) Multilocus genotyping of *P. infestans* isolates from 4000 late blight infection sites over 11 years. This survey revealed that the late blight epidemic that started in 2006 was due to the aggressive 13_A2 isolate that became dominant in only 3 years. The number of surveyed isolates and dominant genotypes of each mating type are indicated. Genotypes with low frequency are grouped as ‘misc’. The shading between bars indicates mating types. (Figure and legend are adapted from Cooke *et al.* [35].) (b) The 13_A2 isolate has rapidly spread to other continents and caused devastating economic losses especially in developing countries such as India and Egypt [38]. Dots represent an approximate of sampling sites for years 2013 and 2014 according to EuroBlight data and recent reports [39].

and symbiosis at the molecular level [46]. Publically available genomic resources are currently limited, which impairs research progress.

5. *Phytophthora ramorum*: the sudden oak death pathogen

Phytophthora ramorum is a devastating oomycete pathogen that causes sudden oak death disease [47]. It has a wide host range, affecting more than 100 plant species from 40 different genera. These include environmentally and economically important tree species such as oaks, tanoaks, rhododendron, European beech, Japanese larch and many woody ornamental plants [47–49]. Disease symptoms are host specific, but

can vary from necrotic lesions in leaves, to shoot dieback and bleeding cankers on the stem [10,50].

In Europe, *P. ramorum* is mainly present in ornamental nurseries or gardens, however, in the UK this pathogen has caused significant tree losses and landscape-scale epidemics [48]. Despite the known reports of *P. ramorum* in Europe [51] and North America [52–55], predictive models suggest that the pathogen might be adapted to larger areas in Africa, Australasia and South America. Therefore, potential of invasion and further spread of the pathogen is possible [56]. In addition, given that diagnosis is primarily based on visual inspection, disease symptoms can be masked by agrochemicals, increasing the likelihood of pathogen intrusion into susceptible areas by the ornamental plant trade market [57].

Phytophthora ramorum occurs as four clonal lineages [58] that have resulted in the emergence of new diseases in forest

ecosystems within Europe and the USA [51,53,59]. The *P. ramorum* genome was sequenced in 2006, a few years after the pathogen was first detected, providing novel insights into its biology [60]. Recent genome sequencing of additional isolates provided further genetic information to track the spread of the pathogen [61]. However, our understanding of its emergence and adaptation remains relatively limited [62].

6. *Plasmopara obducens*: the downy mildew of *Impatiens* species

Plasmopara obducens is an obligate biotrophic pathogen of horticultural plants from the *Impatiens* genus. It causes the *Impatiens* downy mildew foliar disease, which results in wilted and defoliated plants that die within weeks of disease onset [63]. Commercially grown varieties of *Impatiens walleriana* are susceptible to *P. obducens*, making *P. obducens* a clear and present threat to the cultivation of *Impatiens* spp. worldwide [64].

Within the past decade, outbreaks have been reported in North America [65,66], the Hawaiian Islands [67], Europe [68–70], Asia [63] and Australia [71]. Such epidemics were associated with important economic losses [67]. In the USA, the spread of *P. obducens* caused great concern and resulted in the establishment of a research consortium [72]. Cultural practices and chemical applications limit the spread of the disease, however, they are not permanent solutions, as mefenoxam-insensitive *P. obducens* isolates have already been identified [72]. Additionally, *P. obducens* is a persistent pathogen; its ability to spread by wind and survive in soil means disease eradication will be difficult.

The lack of information on the genetics and the structure of *P. obducens* populations, as well as on the factors that led to the establishment of global epidemics, impair the development of effective mitigation strategies. To fill this knowledge gap, Salgado-Salazar *et al.* [28] recently released a draft genome sequence of *P. obducens*, along with a set of genetic markers. These molecular resources may help in determining both the population structure and genetic factors that control infection processes.

7. *Aphanomyces invadans*: the fish epizootic ulcerative syndrome pathogen

Aphanomyces invadans causes epizootic ulcerative syndrome, a fish disease listed on the World Organisation for Animal Health website (<http://www.oie.int/en>). *Aphanomyces invadans* is pathogenic on several economically important fish, including carp, perch, tilapia, snakeheads, salmonids and estuarine fish species [73–75]. It was first reported from farmed ayu, *Plecoglossus altivelis*, in 1971 in Japan [76]. Since then it has spread rapidly throughout Asia and into some areas of Africa.

Aphanomyces invadans has been responsible for large-scale mortalities of farmed and wild fish in more than 20 countries across four continents [77]. In Asia and Africa, the disease has negatively impacted the livelihood of fish farmers and fishermen and, in some cases, threatened the sustainable food supply for local populations, who depend on fish as a relatively affordable source of animal protein [74]. Experimental infection studies have demonstrated that *A. invadans* can produce severe pathological changes in several European salmonid and catfish

species [78]. Hence, it is possible that it could cause serious problems in European tilapia, salmon and trout aquaculture industries and in numerous native and wild fish species.

Despite its economic and social importance, little is known about the biology of the *A. invadans* pathogen. In an effort to address this gap in knowledge, the first draft genome sequence has been completed at the Broad Institute (http://www.ebi.ac.uk/ena/data/view/GCA_000520115). These sequences will enable studies on the biology, evolution, biodiversity, genetics, virulence/pathogenicity and biochemistry of *A. invadans*.

8. *Saprolegnia parasitica*: the saprolegniosis water mould

Saprolegnia parasitica causes saprolegniosis on various fish species. In addition to fish, species of amphibians [79], crustaceans [80] and aquatic insects [81] are also highly susceptible to saprolegniosis. There is conclusive evidence that *Saprolegnia* species are major killers of populations of amphibians globally, threatening some already highly endangered species [82–84]. *Saprolegnia parasitica* is a major problem in the fish farming industry in Europe, Chile, Canada and Asia. At least 10% of all hatched salmon [13] and over 10% of all eggs succumb to *Saprolegnia* infections according to estimates [4,13].

Until 2002, *S. parasitica* was kept under control with applications of malachite green. However, the use of malachite green has been banned worldwide due to its toxicity [85–87], resulting in a dramatic resurgence of *Saprolegnia* infections in salmon aquaculture. At present, fish farmers are struggling to control this pathogen. Current control methods involve treatments with formalin-based products, which are also expected to be banned in the EU in the very near future [88–90]. Therefore, sustainable prevention and control measures are desperately needed.

Recent studies have elucidated an induced native immune response and a suppressed adaptive immune response in fish infected by *S. parasitica* [91]. In addition to distinct biochemical pathways that were described during fish–*Saprolegnia* interaction to establish infection [92], the genome sequence of *S. parasitica* [29,93] has also shed light on the molecular mechanisms of infection, broadening our understanding of infection mechanisms.

9. *Haliotidida noduliformans*: the abalone tubercle mycosis pathogen

Haliotidida noduliformans is a marine pathogen of abalone (*Haliotis* sp.), on which it causes abalone tubercle mycosis, and mantis shrimp (*Oratosquilla oratoria*). It is a member of the Haliphthorales, an early diverged lineage in the Oomycota phylum [94]. *Haliotidida noduliformans* was first discovered in 2004 in South Africa [95,96] and is a threat for abalone in commercial aquaculture, which constitutes more than 93% of the global abalone market [97].

Between 2004 and 2006, several outbreaks in commercial aquaculture farms in South Africa and Japan caused up to 90% mortality among spat and up to 30% mortality among older animals [96,98]. Pathogen control is challenging [96]. Indeed, single *H. noduliformans* isolates can infect at least three abalone species from South Africa, Mexico and Japan [96].

Researchers from Japan and South Africa have provided insights into the distribution of *H. noduliformans* outbreaks and the molecular phylogeny of *Haliotidida* species [95,96,98,99]. Additionally, Greeff *et al.* [100] have established a rapid and sensitive qPCR assay for species-specific detection and quantification of *H. noduliformans* in abalone tissue. However, important resources like genome sequences and gene expression data are lacking. Identification of natural hosts and determining whether these hosts act as reservoirs for *H. noduliformans* will be vital to successfully control this disease.

10. Five ways to tackle emerging or re-emerging oomycete threats

Outbreaks caused by oomycete diseases are a clear and present threat to food security and to natural ecosystems. Although there is a general awareness about these outbreaks, there are gaps in the community capacity to deliver effective short- and long-term response plans. Research showed that oomycetes evolved the ability to infect plants and animals independently of other eukaryotic microbes, and therefore, likely developed unique mechanisms of pathogenicity [101]. The last few decades have seen significant progress in understanding the biology and molecular basis of host infection by oomycete pathogens. Notably, genomic studies have revealed fundamental concepts that link genome architecture, pathogenicity-related proteins and evolution, key factors that drive disease emergence and pathogen success [22]. Here, we highlight five main research areas that need to be sustained or reinforced to help us tackle emerging and re-emerging oomycete threats:

- *Genomic resources.* Sequencing of plant oomycetes has pioneered our understanding of pathogenicity and has provided downstream tools that are essential for mechanistic research. In addition, as more genomes are sequenced, comparative genomic studies will help elucidate infection mechanisms across multiple isolates and species, providing critical information on conserved infection strategies. Oomycete pathogenomics have greatly impacted our approach to plant disease resistance breeding. Indeed, effector proteins have turned out to be useful tools for probing plant germplasm for new resistance traits, and can be used in pathogen-informed resistance breeding programmes. For animal pathogen oomycetes, limited genomic resources are available at the moment (table 1). More efforts are needed for identifying pathogen proteins that manipulate host immune systems, as these will allow for the selection and testing of suitable antigens, which may result in the development of vaccines [102].
- *Genetic manipulation of hosts and pathogens.* Host genetic improvement can be accomplished by precise genome editing techniques, such as the CRISPR/Cas-9 technology [103]. CRISPR/Cas-9 was successfully implemented in tomato to enhance resistance against an oomycete pathogen [104]. To date, mechanistic studies dissecting oomycete–host interactions have been limited to genetic manipulation of the host species due to the lack of efficient gene replacement methods in oomycetes. Recently, Fang & Tyler [105] implemented the CRISPR/Cas9 system in oomycetes and were able to rapidly and efficiently edit the *Phytophthora*

sojae genome. Further implementation of reverse genetic tools in other oomycete species is now needed.

- *Surveillance.* Surveillance consists of the assessment of disease incidence and virulence characterization (via pathotype surveys). In agriculture, pathogen surveillance is crucial to establish national and international policies, government investment and strategies in plant protection, plant breeding, seed distribution and pathology. Surveillance remains a critical aspect of disease control, since understanding the pathogen population allows rapid deployment of control methods. In addition to managed agricultural ecosystems, pathogen surveillance should also include natural habitats and potential reservoirs, as these areas generate new inoculum for re-infection.

We have entered a new era in incursion reporting, where social media provide an efficient and rapid platform for data dissemination and collaborative efforts to tackle new disease emergence and spread. By communicating disease significance to the public and building up awareness of these destructive diseases, a community effort for surveillance can be achieved. This was successfully demonstrated in Australia, after the incursion of myrtle rust [106] and in the recent actions surrounding the outbreak of wheat blast in Bangladesh [107]. Websites such as <http://rusttracker.cimmyt.org> and <http://euroblight.net> are great examples of successful first steps to coordinating global disease surveillance efforts.

- *Field pathogenomics.* In addition to tracking movement and variation of pathogens (surveillance), tackling emerging diseases will not be successful without concurrently understanding the genetics of host resistance. Field pathogenomics adds highly informative data to surveillance surveys by enabling the rapid evaluation of pathogen population structure and host genotype [108]. This has already been successfully implemented with oomycete and fungal pathogens of potato and wheat crops, and will be applicable to other pathogens as more genomic information is generated [35,109].
- *Databases.* Committing to long-term data recording, linkage between databases, free flow of information and faster exchange of information are necessary for updating epidemiological data. Data should be made rapidly and publically available to maximize the combined and coordinated efforts of the scientific community. For instance, scientists recently released websites to share sequencing data generated as an immediate response to outbreaks such as ash dieback (<https://geefu.oadb.tsl.ac.uk/>), wheat yellow rust in the UK (<http://yellowrust.com/>) and wheat blast in Bangladesh (<http://wheatblast.net>). Additionally, social media websites, such as Plant Village (<https://www.plantvillage.org>), allow for free sharing of crop disease information in real time. These websites not only become platforms for scientists to interact, establish collaborations and coordinate among themselves, but also provide information for the general public and farmers to directly follow and participate in research efforts. They allow the world to connect, share and collectively find solutions to emerging oomycete threats.

Fighting agriculturally important pathogens is a long-term ongoing battle. Just like the constantly morphing flu virus, oomycete pathogens continuously evolve new races to evade host resistance. Agricultural scientists, veterinarians,

breeders and farmers need to remain vigilant and work together in reporting and tackling oomycete pathogens to achieve successful control.

Authors' contributions. S.K. outlined the article. All authors were assigned sections of the article and drafted the associated text. L.D. and S.K. oversaw the revisions.

Competing interests. We have no competing interests.

Funding. The Kamoun Lab is funded by the Biotechnology and Biological Sciences Research Council, the European Research Council

and the Gatsby Charitable Foundation. R.K. is funded by CEPLAS—Cluster of Excellence on Plant Sciences (EXC 1028). J.C.D.I.C. is funded by The John Innes Foundation. P.v.W. is funded by the Biotechnology and Biological Sciences Research Council, the Natural Environment Research Council, and the Newton Fund.

Acknowledgements. We thank Doug Caldwell for inspiring us to select *P. obducens*. The authors would also like to thank Jimmy Wales and Larry Sanger for providing a platform for dissemination of useful knowledge.

References

- Baldauf SL, Roger AJ, Wenk-Siefert I, Doolittle WF. 2000 A kingdom-level phylogeny of eukaryotes based on combined protein data. *Science* **290**, 972–977. (doi:10.1126/science.290.5493.972)
- Beakes GW, Glockling SL, Sekimoto S. 2012 The evolutionary phylogeny of the oomycete 'fungi'. *Protospasma* **249**, 3–19. (doi:10.1007/s00709-011-0269-2)
- Dieguez-Urbeondo J *et al.* 2009 Phylogenetic relationships among plant and animal parasites, and saprotrophs in *Aphanomyces* (Oomycetes). *Fungal Genet. Biol.* **46**, 365–376. (doi:10.1016/j.fgb.2009.02.004)
- van West P. 2006 *Saprolegnia parasitica*, an oomycete pathogen with a fishy appetite: new challenges for an old problem. *Mycologist* **20**, 99–104. (doi:10.1016/j.mycol.2006.06.004)
- Kamoun S, Smart CD. 2005 Late blight of potato and tomato in the genomics era. *Plant Dis.* **89**, 692–699. (doi:10.1094/PD-89-0692)
- Kamoun S. 2003 Molecular genetics of pathogenic oomycetes. *Eukaryot. Cell* **2**, 191–199. (doi: 10.1128/EC.2.2.191-199.2003)
- Erwin DC, Ribeiro OK. 1996 *Phytophthora diseases worldwide*. St. Paul, MN: American Phytopathology Society Press.
- Haas BJ *et al.* 2009 Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* **461**, 393–398. (doi:10.1038/nature08358)
- Drenth A, Guest DL. 2013 *Phytophthora palmivora* in tropical tree crops. In *Phytophthora: a global perspective* (ed. K Lamour), p. 187. Wallingford, UK: CAB.
- Rizzo DM, Garbelotto M, Hansen EM. 2005 *Phytophthora ramorum*: integrative research and management of an emerging pathogen in California and Oregon forests. *Annu. Rev. Phytopathol.* **43**, 309–335. (doi:10.1146/annurev.phyto.42.040803.140418)
- Agrios GN. 2004 *Plant pathology*. 5th edn. San Diego, CA: Academic.
- Lévesque CA *et al.* 2010 Genome sequence of the necrotrophic plant pathogen *Pythium ultimum* reveals original pathogenicity mechanisms and effector repertoire. *Genome Biol.* **11**, 1–22. (doi:10.1186/gb-2010-11-7-r73)
- Phillips AJ, Anderson VL, Robertson EJ, Secombes CJ, van West P. 2008 New insights into animal pathogenic oomycetes. *Trends Microbiol.* **16**, 13–19. (doi:10.1016/j.tim.2007.10.013)
- FAO. 2012 *The state of world fisheries and aquaculture 2012*. Rome, Italy: FAO.
- Martini R, Lindberg C. 2013 *Fishing for tomorrow: managing fisheries for sustainable development*. Organisation for Economic Co-operation and Development. https://www.oecd.org/pcd/002013401_Coherence%20for%20Development_Dec_PDFX_WEB.pdf.
- van West P, Beakes GW. 2014 Animal pathogenic oomycetes. *Fungal Biol.* **118**, 525–526. (doi:10.1016/j.funbio.2014.05.004)
- Thines M. 2014 Phylogeny and evolution of plant pathogenic oomycetes—a global overview. *Eur. J. Plant Pathol.* **138**, 431–447. (doi:10.1007/s10658-013-0366-5)
- McDonald BA, Linde C. 2002 Pathogen population genetics, evolutionary potential, and durable resistance. *Annu. Rev. Phytopathol.* **40**, 349–379. (doi:10.1146/annurev.phyto.40.120501.101443)
- Bozkurt TO, Schornack S, Banfield MJ, Kamoun S. 2012 Oomycetes, effectors, and all that jazz. *Curr. Opin. Plant Biol.* **15**, 483–492. (doi:10.1016/j.pbi.2012.03.008)
- Schornack S *et al.* 2009 Ten things to know about oomycete effectors. *Mol. Plant Pathol.* **10**, 795–803. (doi:10.1111/j.1364-3703.2009.00593.x)
- Raffaele S, Kamoun S. 2012 Genome evolution in filamentous plant pathogens: why bigger can be better. *Nat. Rev. Microbiol.* **10**, 417–430. (doi:10.1038/nrmicro2790)
- Dong S, Raffaele S, Kamoun S. 2015 The two-speed genomes of filamentous pathogens: waltz with plants. *Curr. Opin. Genet. Dev.* **35**, 57–65. (doi:10.1016/j.gde.2015.09.001)
- Daggett SS, Knighton JE, Therrien CD. 1995 Polyploidy among isolates of *Phytophthora infestans* from Eastern Germany. *J. Phytopathol.* **143**, 419–422. (doi:10.1111/j.1439-0434.1995.tb00286.x)
- Yoshida K *et al.* 2013 The rise and fall of the *Phytophthora infestans* lineage that triggered the Irish potato famine. *eLife* **2**, e00731. (doi:10.7554/eLife.00731)
- Hamed BH, Gisi U. 2013 Generation of pathogenic F1 progeny from crosses of *Phytophthora infestans* isolates differing in ploidy. *Plant Pathol.* **62**, 708–718. (doi:10.1111/j.1365-3059.2012.02655.x)
- Li Y, Zhou Q, Qian K, van der Lee T, Huang SW. 2015 Successful asexual lineages of the Irish potato famine pathogen are triploid. *bioRxiv* 024596. (doi:10.1101/024596)
- Tyler BM *et al.* 2006 *Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis. *Science* **313**, 1261–1266. (doi:10.1126/science.1128796)
- Salgado-Salazar C, Rivera Y, Veltri D, Crouch JA. 2015 Polymorphic SSR markers for *Plasmopara obducens* (Peronosporaceae), the newly emergent downy mildew pathogen of *Impatiens* (Balsaminaceae). *Appl. Plant Sci.* **3**, 1500073. (doi:10.3732/apps.1500073)
- Jiang RH *et al.* 2013 Distinctive expansion of potential virulence genes in the genome of the oomycete fish pathogen *Saprolegnia parasitica*. *PLoS Genet.* **9**, e1003272. (doi:10.1371/journal.pgen.1003272)
- Haverkort AJ, Boonekamp PM, Hutten R, Jacobsen E, Lotz LAP, Kessel GJT, Visser RGF, van der Vossen EAG. 2008 Societal costs of late blight in potato and prospects of durable resistance through cisgenic modification. *Potato Res.* **51**, 47–57. (doi:10.1007/s11540-008-9089-y)
- Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012 Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**, 186–194. (doi:10.1038/nature10947)
- Fry WE *et al.* 2015 Five reasons to consider *Phytophthora infestans* a reemerging pathogen. *Phytopathology* **105**, 966–981. (doi:10.1094/PHYTO-01-15-0005-FI)
- Kato M, Mizubuti ES, Goodwin SB, Fry WE. 1997 Sensitivity to protectant fungicides and pathogenic fitness of clonal lineages of *Phytophthora infestans* in the United States. *Phytopathology* **87**, 973–978. (doi:10.1094/PHYTO.1997.87.9.973)
- Fry W. 2008 *Phytophthora infestans*: the plant (and R gene) destroyer. *Mol. Plant Pathol.* **9**, 385–402. (doi:10.1111/j.1364-3703.2007.00465.x)
- Cooke DE *et al.* 2012 Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathog.* **8**, e1002940. (doi:10.1371/journal.ppat.1002940)
- Fry WE *et al.* 2013 The 2009 late blight pandemic in the Eastern United States—causes and results.

- Plant Dis.* **97**, 296–306. (doi:10.1094/pdis-08-12-0791-fe)
37. Harbaoui K, Hamada W, Li Y, Vleeshouwers VGAA, van der Lee T. 2014 Increased difficulties to control late blight in Tunisia are caused by a genetically diverse *Phytophthora infestans* population next to the clonal lineage NA-01. *Plant Dis.* **98**, 898–908. (doi:10.1094/pdis-06-13-0610-re)
38. Chowdappa P, Kumar NBJ, Madhura S, Kumar MSP, Myers KL, Fry WE, Squires JN, Cooke DEL. 2013 Emergence of 13_A2 blue lineage of *Phytophthora infestans* was responsible for severe outbreaks of late blight on tomato in South-West India. *J. Phytopathol.* **161**, 49–58. (doi:10.1111/jph.12031)
39. El-Ganainy SM, Tohamy AMM, Awad MA, Squires JN, Cooke DE. 2013 The population dynamics of *Phytophthora infestans* in Egypt. In *American Phytopathological Society Annual Meeting 2013, Austin, TX, 10–14 August*.
40. Raffaele S, Win J, Cano LM, Kamoun S. 2010 Analyses of genome architecture and gene expression reveal novel candidate virulence factors in the secretome of *Phytophthora infestans*. *BMC Genomics* **11**, 637. (doi:10.1186/1471-2164-11-637)
41. Stassen JH, van den Ackerveken G. 2011 How do oomycete effectors interfere with plant life? *Curr. Opin. Plant Biol.* **14**, 407–414. (doi:10.1016/j.pbi.2011.05.002)
42. Torres GA, Sarria GA, Martinez G, Varon A, Drenth A, Guest DI. 2016 Bud rot caused by *Phytophthora palmivora*: a destructive emerging disease of oil palm. *Phytopathology* **106**, 320–329. (doi:10.1094/PHYTO-09-15-0243-RVW)
43. Rey T, Chatterjee A, Buttany M, Toulotte J, Schornack S. 2015 *Medicago truncatula* symbiosis mutants affected in the interaction with a biotrophic root pathogen. *New Phytol.* **206**, 497–500. (doi:10.1111/nph.13233)
44. Le Fevre R, O'Boyle B, Moscou MJ, Schornack S. 2016 Colonization of barley by the broad-host hemibiotrophic pathogen *Phytophthora palmivora* uncovers a leaf development dependent involvement of MLO. *Mol. Plant Microbe Interact.* **29**, 385–395. (doi:10.1094/MPMI-12-15-0276-R)
45. Chaparro-García A, Wilkinson RC, Gimenez-Ibanez S, Findlay K, Coffey MD, Zipfel C, Rathjen JP, Kamoun S, Schornack S. 2011 The receptor-like kinase SERK3/BAK1 is required for basal resistance against the late blight pathogen *Phytophthora infestans* in *Nicotiana benthamiana*. *PLoS ONE* **6**, e16608. (doi:10.1371/journal.pone.0016608)
46. Wang E, Schornack S, Marsh JF, Gobbato E, Schwessinger B, Eastmond P, Schultze M, Kamoun S, Oldroyd GE. 2012 A common signaling process that promotes mycorrhizal and oomycete colonization of plants. *Curr. Biol.* **22**, 2242–2246. (doi:10.1016/j.cub.2012.09.043)
47. Grünwald NJ, Goss EM, Press CM. 2008 *Phytophthora ramorum*: a pathogen with a remarkably wide host range causing sudden oak death on oaks and ramorum blight on woody ornamentals. *Mol. Plant Pathol.* **9**, 729–740. (doi:10.1111/j.1364-3703.2008.00500.x)
48. Brasier C, Webber J. 2010 Plant pathology: sudden larch death. *Nature* **466**, 824–825. (doi:10.1038/466824a)
49. Grünwald NJ, Garbelotto M, Goss EM, Heungens K, Prospero S. 2012 Emergence of the sudden oak death pathogen *Phytophthora ramorum*. *Trends Microbiol.* **20**, 131–138. (doi:10.1016/j.tim.2011.12.006)
50. Hansen EM, Parke JL, Sutton W. 2005 Susceptibility of Oregon forest trees and shrubs to *Phytophthora ramorum*: a comparison of artificial inoculation and natural infection. *Plant Dis.* **89**, 63–70. (doi:10.1094/pd-89-0063)
51. Vercauteren A, Dobbelaere I, Bockstaele E, Maes M, Heungens K. 2010 Genotypic and phenotypic characterization of the European A2 isolates of *Phytophthora ramorum*. *Eur. J. Plant Pathol.* **129**, 621–635. (doi:10.1007/s10658-010-9727-5)
52. Goss EM, Carbone I, Grünwald NJ. 2009 Ancient isolation and independent evolution of the three clonal lineages of the exotic sudden oak death pathogen *Phytophthora ramorum*. *Mol. Ecol.* **18**, 1161–1174. (doi:10.1111/j.1365-294X.2009.04089.x)
53. Goss EM, Larsen M, Chastagner GA, Givens DR, Grünwald NJ. 2009 Population genetic analysis infers migration pathways of *Phytophthora ramorum* in US nurseries. *PLoS Pathog.* **5**, e1000583. (doi:10.1371/journal.ppat.1000583)
54. Ivors K, Garbelotto M, Vries ID, Ruyter-Spira C, Te Hekkert B, Rosenzweig N, Bonants P. 2006 Microsatellite markers identify three lineages of *Phytophthora ramorum* in US nurseries, yet single lineages in US forest and European nursery populations. *Mol. Ecol.* **15**, 1493–1505. (doi:10.1111/j.1365-294X.2006.02864.x)
55. Mascheretti S, Croucher PJ, Kozanits M, Baker L, Garbelotto M. 2009 Genetic epidemiology of the sudden oak death pathogen *Phytophthora ramorum* in California. *Mol. Ecol.* **18**, 4577–4590. (doi:10.1111/j.1365-294X.2009.04379.x)
56. Ireland KB, Hardy GE, Kriticos DJ. 2013 Combining inferential and deductive approaches to estimate the potential geographical range of the invasive plant pathogen, *Phytophthora ramorum*. *PLoS ONE* **8**, e63508. (doi:10.1371/journal.pone.0063508)
57. Moslonka-Lefebvre M, Finlay A, Dorigatti I, Dehnen-Schmutz K, Harwood T, Jeger MJ, Xu X, Holdenrieder O, Pautasso M. 2011 Networks in plant epidemiology: from genes to landscapes, countries, and continents. *Phytopathology* **101**, 392–403. (doi:10.1094/PHYTO-07-10-0192)
58. Van Poucke K, Franceschini S, Webber JF, Vercauteren A, Turner JA, McCracken AR, Heungens K, Brasier CM. 2012 Discovery of a fourth evolutionary lineage of *Phytophthora ramorum*: EU2. *Fungal Biol.* **116**, 1178–1191. (doi:10.1016/j.funbio.2012.09.003)
59. Goss EM, Larsen M, Vercauteren A, Werres S, Heungens K, Grünwald NJ. 2010 *Phytophthora ramorum* in Canada: evidence for migration within North America and from Europe. *Phytopathology* **101**, 166–171. (doi:10.1094/PHYTO-05-10-0133)
60. Kamoun S *et al.* 2015 The top 10 oomycete pathogens in molecular plant pathology. *Mol. Plant Pathol.* **16**, 413–434. (doi:10.1111/mpp.12190)
61. de la Mata Saez L, McCracken AR, Cooke LR, O'Neill P, Grant M, Studholme DJ. 2015 Draft genome sequences of seven isolates of *Phytophthora ramorum* EU2 from Northern Ireland. *Genom. Data* **6**, 191–192. (doi:10.1016/j.gdata.2015.09.009)
62. Grünwald NJ. 2012 Genome sequences of *Phytophthora* enable translational plant disease management and accelerate research. *Can. J. Plant Pathol. Rev. Can. Phytopathol.* **34**, 13–19. (doi:10.1080/07060661.2012.664568)
63. Shen YM, Huang JH, Liu HL. 2013 First report of downy mildew caused by *Plasmopara obducens* on impatiens in Taiwan. *Plant Dis.* **97**, 1512. (doi:10.1094/PDIS-05-13-0501-PDN)
64. Braiser CM. 2008 The biosecurity threat to the UK and global environment from international trade in plants. *Plant Pathol.* **57**, 792–808. (doi:10.1111/j.1365-3059.2008.01886.x)
65. Wegulo SN, Koike ST, Vilchez M, Santos P. 2004 First report of downy mildew caused by *Plasmopara obducens* on impatiens in California. *Plant Dis.* **88**, 909. (doi:10.1094/PDIS.2004.88.8.909B)
66. Baysal-Gurel F, Taylor NJ, Chatfield J, Miller SA. 2012 First report of impatiens downy mildew caused by *Plasmopara obducens* in Ohio. *Plant Dis.* **96**, 1699. (doi:10.1094/PDIS-06-12-0545-PDN)
67. Crouch JA, Ko MP, McKemy JM. 2014 First report of impatiens downy mildew outbreaks caused by *Plasmopara obducens* throughout the Hawaiian islands. *Plant Dis.* **98**, 696. (doi:10.1094/PDIS-10-13-1017-PDN)
68. Lane CR, Beales PA, O'Neill TM, McPherson GM, Finlay AR, David J, Constantinescu O, Henricot B. 2005 First report of *Impatiens* downy mildew (*Plasmopara obducens*) in the UK. *Plant Pathol.* **54**, 243. (doi:10.1111/j.1365-3059.2005.01133.x)
69. Toppe B, Brurberg MB, Stensvand A, Herrero ML. 2010 First report of *Plasmopara obducens* (downy mildew) on *Impatiens walleriana* in Norway. *Plant Pathol.* **59**, 800. (doi:10.1111/j.1365-3059.2010.02270.x)
70. Bulajic A, Vucurovic A, Stankovic I, Ristic D, Jovic J, Stojkovic B, Krstic B. 2011 First report of *Plasmopara obducens* on *Impatiens walleriana* in Serbia. *Plant Dis.* **95**, 491. (doi:10.1094/PDIS-10-10-0731)
71. Cunnington JH, Aldaoud R, Loh M, Washington WS, Irvine G. 2008 First record of *Plasmopara obducens* (downy mildew) on impatiens in Australia. *Plant Pathol.* **57**, 371. (doi:10.1111/j.1365-3059.2007.01630.x)
72. Daughtry M, Cristi P. 2014 Impatiens: is there life after downy mildew? *American Hort News. American Hort News* 11 March.
73. Lilley JH, Hart D, Panyawachira V, Kanchanakhan S, Chinabut S, Soderhall K, Cerenius L. 2003 Molecular characterization of the fish-pathogenic fungus *Aphanomyces invadans*. *J. Fish Dis.* **26**, 263–275. (doi:10.1046/j.1365-2761.2003.00448.x)

74. Oidtman B. 2012 Review of biological factors relevant to import risk assessments for epizootic ulcerative syndrome (*Aphanomyces invadans*). *Transbound. Emerg. Dis.* **59**, 26–39. (doi:10.1111/j.1865-1682.2011.01241.x)
75. Vandersea MW *et al.* 2006 Molecular assays for detecting *Aphanomyces invadans* in ulcerative mycotic fish lesions. *Appl. Environ. Microbiol.* **72**, 1551–1557. (doi:10.1128/AEM.72.2.1551-1557.2006)
76. Egusa S, Masuda N. 1971 [A new fungal disease of *Plecoglossus altivelis*]. *Fish Pathol.* **6**, 41–46. (doi:10.3147/jstfp.6.41) [In Japanese.]
77. Pradhan PK *et al.* 2014 Emergence of epizootic ulcerative syndrome: large-scale mortalities of cultured and wild fish species in Uttar Pradesh, India. *Curr. Sci.* **106**, 1711–1718.
78. Oidtman B, Steinbauer P, Geiger S, Hoffmann RW. 2008 Experimental infection and detection of *Aphanomyces invadans* in European catfish, rainbow trout and European eel. *Dis. Aquat. Organ.* **82**, 195–207. (doi:10.3354/dao01973)
79. Blaustein AR, Hokit DG, O'Hara RK, Holt RA. 1994 Pathogenic fungus contributes to amphibian losses in the pacific northwest. *Biol. Conserv.* **67**, 251–254. (doi:10.1016/0006-3207(94)90616-5)
80. Dieguez-Urbeondo J, Cerenius L, Soderhall K. 1994 *Saprolegnia parasitica* and its virulence on three different species of freshwater crayfish. *Aquaculture* **120**, 219–228. (doi:10.1016/0044-8486(94)90080-9)
81. Sarowar MN, van den Berg AH, McLaggan D, Young MR, van West P. 2013 *Saprolegnia* strains isolated from river insects and amphipods are broad spectrum pathogens. *Fungal Biol.* **117**, 752–763. (doi:10.1016/j.funbio.2013.09.002)
82. Fernández-Benítez MJ, Ortiz-Santaliestra ME, Lizana M, Diéguez-Urbeondo J. 2011 Differences in susceptibility to *Saprolegnia* infections among embryonic stages of two anuran species. *Oecologia* **165**, 819–826. (doi:10.1007/s00442-010-1889-5)
83. Fernández-Benítez MJ, Ortiz-Santaliestra ME, Lizana M, Diéguez-Urbeondo J. 2008 *Saprolegnia diclina*: another species responsible for the emergent disease 'Saprolegnia infections' in amphibians. *FEMS Microbiol. Lett.* **279**, 23–29. (doi:10.1111/j.1574-6968.2007.01002.x)
84. Kiesecker JM, Blaustein AR, Miller CL. 2001 Transfer of a pathogen from fish to amphibians. *Conserv. Biol.* **15**, 1064–1070. (doi:10.1046/j.1523-1739.2001.0150041064.x)
85. Alderman DJ, Polglase JL. 1984 A comparative investigation of the effects of fungicides on *Saprolegnia parasitica* and *Aphanomyces astaci*. *Trans. Br. Mycol. Soc.* **83**, 313–318. (doi:10.1016/S0007-1536(84)80153-9)
86. Meyer FP, Jorgenson TA. 1983 Teratological and other effects of malachite green on the development of rainbow trout and rabbits. *Trans. Am. Fish Soc.* **112**, 818–824. (doi:10.1577/1548-8659(1983)112<818:TAOEOEOM>2.0.CO;2)
87. Srivastava S, Sinha R, Roy D. 2004 Toxicological effects of malachite green. *Aquat. Toxicol.* **66**, 319–329. (doi:10.1016/j.aquatox.2003.09.008)
88. Magaraggia M, Faccenda F, Gandolfi A, Jori G. 2006 Treatment of microbiologically polluted aquaculture waters by a novel photochemical technique of potentially low environmental impact. *J. Environ. Monit.* **8**, 923–931. (doi:10.1039/B606975D)
89. Marking LL, Rach JJ, Schreier TM. 1994 Evaluation of antifungal agents for fish culture. *Progress. Fish Culturist* **56**, 225–231. (doi:10.1577/1548-8640(1994)056<0225:AFSEOA>2.3.CO;2)
90. Burka JF, Hammell KL, Horsberg TE, Johnson GR, Rainnie DJ, Speare DJ. 1997 Drugs in salmonid aquaculture—a review. *J. Vet. Pharmacol. Ther.* **20**, 333–349. (doi:10.1046/j.1365-2885.1997.00094.x)
91. Belmonte R, Wang T, Duncan GJ, Skaar I, Mélida H, Bulone V, van West P, Secombes CJ. 2014 Role of pathogen-derived cell wall carbohydrates and prostaglandin E2 in immune response and suppression of fish immunity by the oomycete *Saprolegnia parasitica*. *Infect. Immun.* **82**, 4518–4529. (doi:10.1128/IAI.02196-14)
92. Wawra S *et al.* 2012 Host-targeting protein 1 (SpHtp1) from the oomycete *Saprolegnia parasitica* translocates specifically into fish cells in a tyrosine-O-sulphate-dependent manner. *Proc. Natl Acad. Sci. USA* **109**, 2096–2101. (doi:10.1073/pnas.1113775109)
93. Wawra S, Belmonte R, Löblich L, Saraiva M, Willems A, van West P. 2012 Secretion, delivery and function of oomycete effector proteins. *Curr. Opin. Microbiol.* **15**, 685–691. (doi:10.1016/j.mib.2012.10.008)
94. Beakes D, Honda D, Thines M. 2014 Systematics of the Straminipila: Labyrinthulomycota, Hyphochytriomycota and Oomycota. In *The Mycota—systematics and evolution part A*, 2nd edn (eds DJ McLaughlin, JW Spatafora), pp. 39–97. Berlin, Germany: Springer.
95. Atami H, Muraosa Y, Hatai K. 2009 *Halioticida* infection found in wild mantis shrimp *Oratosquilla oratoria* in Japan. *Fish Pathol.* **44**, 145–150. (doi:10.3147/jstfp.44.145)
96. Muraosa Y, Morimoto K, Sano A, Nishimura K, Hatai K. 2009 A new peronosporomycete, *Halioticida noduliformans* gen. et sp. nov., isolated from white nodules in the abalone *Haliotis* spp. from Japan. *Mycoscience* **50**, 106–115. (doi:10.1007/s10267-008-0462-0)
97. Cook PA. 2014 The worldwide abalone industry. *Mod. Econ.* **5**, 1181–1186. (doi:10.4236/me.2014.513110)
98. Macey BM, Christison KW, Mouton A. 2011 *Halioticida noduliformans* isolated from cultured abalone (*Haliotis midae*) in South Africa. *Aquaculture* **315**, 187–195. (doi:10.1016/j.aquaculture.2011.02.004)
99. Hatai K. 2012 Diseases of fish and shellfish caused by marine fungi. In *Biology of marine fungi* (ed. C Rsaghukumar), pp. 159–171. Berlin, Germany: Springer.
100. Greeff MR, Christison KW, MacEay BM. 2012 Development and preliminary evaluation of a real-time PCR assay for *Halioticida noduliformans* in abalone tissues. *Dis. Aquat. Organ.* **99**, 103–117. (doi:10.3354/dao02468)
101. Jiang RHY, Tyler BM. 2012 Mechanisms and evolution of virulence in oomycetes. *Annu. Rev. Phytopathol.* **50**, 295–318. (doi:10.1146/annurev-phyto-081211-172912)
102. Minor K *et al.* 2014 A putative serine protease, SpSsp1, from *Saprolegnia parasitica* is recognised by sera of rainbow trout, *Oncorhynchus mykiss*. *Fungal Biol.* **118**, 630–639. (doi:10.1016/j.funbio.2014.04.008)
103. Belhaj K, Chaparro-García A, Kamoun S, Patron NJ, Nekrasov V. 2014 Engineering plant genomes with CRISPR/Cas9. *Curr. Opin. Biotechnol.* **32**, 76–84. (doi:10.1016/j.copbio.2014.11.007)
104. de Toledo Thomazella DP, Brail Q, Dahlbeck D, Staskawicz BJ. 2016 CRISPR-Cas9 mediated mutagenesis of a DMR6 ortholog in tomato confers broad-spectrum disease resistance. *bioRxiv*. (doi:10.1101/064824)
105. Fang Y, Tyler BM. 2016 Efficient disruption and replacement of an effector gene in the oomycete *Phytophthora sojae* using CRISPR/cas9. *Mol. Plant Pathol.* **17**, 127–139. (doi:10.1111/mpp.12318)
106. Carnegie AJ, Cooper K. 2011 Emergency response to the incursion of an exotic myrtaceous rust in Australia. *Australas. Plant Pathol.* **40**, 346. (doi:10.1007/s13313-011-0066-6)
107. Islam T *et al.* 2016 Emergence of wheat blast in Bangladesh was caused by a South American lineage of *Magnaporthe oryzae*. *bioRxiv*. (doi:10.1101/059832)
108. Derevnina L, Michelmore RW. 2015 Wheat rust never sleep but neither do sequencers: will pathogenics transform the way plant disease is managed? *Genome Biol.* **16**, 44. (doi:10.1186/s13059-015-0615-3)
109. Hubbard A *et al.* 2015 Field pathogenomics reveals the emergence of a diverse wheat yellow rust population. *Genome Biol.* **16**, 23. (doi:10.1186/s13059-015-0590-8)