

Sequence-Based Species Delimitation for the DNA Taxonomy of Undescribed Insects

JOAN PONS,^{1,2,10} TIMOTHY G. BARRACLOUGH,^{2,3} JESUS GOMEZ-ZURITA,^{1,2,7} ANABELA CARDOSO,^{1,4,7}
DANIEL P. DURAN,^{1,8} STEAPHAN HAZELL,^{1,2,9} SOPHIEN KAMOUN,⁵ WILLIAM D. SUMLIN,⁶
AND ALFRIED P. VOGLER^{1,2}

¹Department of Entomology, The Natural History Museum, London SW7 5BD, United Kingdom; E-mail: a.vogler@nhm.ac.uk (A.P.V.)

²Division of Biology and NERC Centre for Population Biology, Imperial College London, Silwood Park Campus, Ascot, Berkshire SL5 7PY, United Kingdom

³Jodrell Laboratory, Royal Botanic Gardens, Kew TW9 3DS, United Kingdom

⁴Faculdade de Ciências da Universidade de Lisboa, Departamento de Biologia Animal, Centro de Biologia Ambiental, Rua Ernesto Vasconcelos, 1749-016, Campo Grande, Lisboa, Portugal

⁵Department of Plant Pathology, Ohio State University, Ohio Agricultural Research and Development Center, Wooster, Ohio, 44691, USA

⁶Department of Entomology, Texas A&M University, College Station, Texas 77843, USA

⁷Present Address: Area de Biología Animal, Departamento de Zoología y Antropología Física, Facultad de Biología, Universidad de Murcia—Campus de Espinardo, 30071 Murcia, Spain

⁸Present Address: Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee, 37235, USA

⁹Present Address: Division of Zoology, School of Animal and Microbial Sciences, University of Reading, Reading RG6 6AJ, United Kingdom

¹⁰Present Address: Unitat de Biologia Evolutiva, Facultat de Ciències de la Salut i e de la Vida, Universitat Pompeu Fabra, C/Dr. Aiguader 80, 08003 Barcelona, Catalonia, Spain

Abstract.— Cataloging the very large number of undescribed species of insects could be greatly accelerated by automated DNA based approaches, but procedures for large-scale species discovery from sequence data are currently lacking. Here, we use mitochondrial DNA variation to delimit species in a poorly known beetle radiation in the genus *Rivacindela* from arid Australia. Among 468 individuals sampled from 65 sites and multiple morphologically distinguishable types, sequence variation in three mtDNA genes (cytochrome oxidase subunit 1, cytochrome *b*, 16S ribosomal RNA) was strongly partitioned between 46 or 47 putative species identified with quantitative methods of species recognition based on fixed unique (“diagnostic”) characters. The boundaries between groups were also recognizable from a striking increase in branching rate in clock-constrained calibrated trees. Models of stochastic lineage growth (Yule models) were combined with coalescence theory to develop a new likelihood method that determines the point of transition from species-level (speciation and extinction) to population-level (coalescence) evolutionary processes. Fitting the location of the switches from speciation to coalescent nodes on the ultrametric tree of *Rivacindela* produced a transition in branching rate occurring at 0.43 Mya, leading to an estimate of 48 putative species (confidence interval for the threshold ranging from 47 to 51 clusters within 2 log L units). Entities delimited in this way exhibited biological properties of traditionally defined species, showing coherence of geographic ranges, broad congruence with morphologically recognized species, and levels of sequence divergence typical for closely related species of insects. The finding of discontinuous evolutionary groupings that are readily apparent in patterns of sequence variation permits largely automated species delineation from DNA surveys of local communities as a scaffold for taxonomy in this poorly known insect group. [Phylogenetic species concept; coalescence; mtDNA; Cicindelidae; Australia; paleoclimate.]

Automated DNA sequencing procedures could greatly accelerate identification of known species (“DNA barcoding”; Hebert et al., 2003) and provide a framework for the taxonomy of poorly known groups (Tautz et al., 2003). However, sequencing-based approaches in taxonomy are useful only if an individual’s DNA sequence can unambiguously associate that individual with a larger group of organisms (a species or higher taxon). Establishing this association is complicated due to intraspecific variation and the possible incongruence of gene and species histories that might prevent the easy recognition of these groups, in particular where only a single fragment of mtDNA is used (Lipscomb et al., 2003; Mallet and Willmott, 2003; Will and Rubinoff, 2004). In current taxonomic DNA sequencing, species-level entities have been accepted to be known a priori, and their DNA signature is based on genotypes from a representative sample of well-identified individuals (Hajibabaei et al., 2006; Hebert et al., 2003; Meyer and Paulay, 2005).

“DNA barcoding” therefore accepts that species limits are established following traditional practices of taxonomy, usually based on morphology, and the DNA information is fitted into this system of predefined taxonomic

groups. This approach is problematic because the correspondence of sequence variation with existing Linnean binomials is expected to be inexact. This may be because of the existence of unrecognized cryptic species (Hebert et al., 2004a, 2004b) or incongruence of gene histories confounding species recognition (Funk and Omland, 2003; Meyer and Paulay, 2005; Monaghan et al., 2006) or because procedures for species delimitation of the traditional taxonomy have been vague or species concepts have been applied inconsistently (Agapow et al., 2004; Cracraft, 1992; Meyer and Paulay, 2005; Wiens and Penkrot, 2002). To avoid these problems, a DNA-based taxonomic system should use the sequence information itself as the primary information source for establishing group membership and defining species boundaries.

Current procedures for delimiting species from sequence data aggregate populations lacking discrete differences into a single species (Cracraft, 1983; Davis and Nixon, 1992; Sites and Marshall, 2003; Wiens and Penkrot, 2002). These “smallest detected sample[s] of self-perpetuating organisms that have unique sets of characters” (Nelson and Platnick, 1981) can be recognized based on fixed nucleotide differences unique to

such (sets of) populations (the criterion of “diagnosability”; Cracraft, 1983), implemented in population aggregation analysis (PAA; Davis and Nixon, 1992). A tree-based variant of this procedure (cladistic haplotype analysis; CHA) also considers homoplastic character states as diagnostic if they are shown to be uniquely derived for a particular group (Brower, 1999). The Wiens-Penkrot (WP) method (Sites and Marshall, 2003; Wiens and Penkrot, 2002) is also a tree-based method for delimiting phylogenetic species but uses the topology (rather than character polarity, as in CHA) to assess whether or not clades are restricted to a (set of) populations to the exclusion of clades elsewhere (criterion of “exclusivity”).

The prior definition of populations, usually defined geographically, is a critical step in all of these methods, although the focus on individual organisms rather than a preconceived interbreeding “group” would be more objective for species delimitation (Vrana and Wheeler, 1992). Current methods for estimating population coherence from the sequences themselves rely on evidence for recombination to determine species limits (Doyle, 1995) but are impractical for large-scale analysis at present. An alternative possibility to detect species separation is to search for discontinuities in sequence variation associated with the species boundaries. For example, statistical parsimony analysis (Templeton et al., 1992) separates groups of sequences into different sequence networks if genotypes are connected by comparatively long branches that are affected by homoplasy. Although homoplasious connections do not necessarily correspond to species boundaries, in practice this algorithm often does separate groups that are roughly coincident with named species or species groups (e.g., Cardoso and Vogler, 2005; Templeton, 2001; Wilder and Hollocher, 2003; Templeton et al., 1995), and the WP method for species delimitation is now recommended to be applied only in conjunction with statistical parsimony analysis in order to reduce the complexity of large data sets (Sites and Marshall, 2003).

These observations would suggest the possibility of using analyses of branch lengths on a DNA tree for explicit tests of species boundaries, based on the difference in branching rates at the level of species and populations. Branch lengths between species are determined by speciation and extinction rates (macroevolution) (Nee et al., 1994), whereas branch lengths within a species reflect coalescence processes at the level of populations (microevolution) (Hudson, 1991; Rosenberg and Nordborg, 2002; Wakeley, 2006). Well-developed approaches exist for analyzing branching rates in either framework. Combining equations that describe processes of lineage birth at the species level with coalescence models within species, it is possible to develop a statistical framework for estimating the predicted shift in dynamics of branching associated with the species boundary. Here we develop a method that determines the locations of ancestral nodes that define putative species and applies a likelihood ratio test to assess the fit of the branch lengths to a mixed lineage birth-population coalescence model.

As an example of a group whose current taxonomy is incomplete, we attempted DNA-based species

delineation in tiger beetles in the genus *Rivacindela* (Coleoptera: Cicindelidae). These beetles are found in saline habitats and temporal lakes of interior Australia. To date, 24 species of *Rivacindela* have been formally described but because of the inaccessibility of their habitat, many species still await discovery (Freitag, 1979; Sumlin, 1997). Populations are restricted to habitat near the water edges in an otherwise inhospitable desert environment. Although similar in their lifestyle as general predators of small arthropods, species and populations differ in color pattern, body shape, habitat preference, and behavioral traits. Most species are flighted, but *Rivacindela* includes several flightless lineages that are extremely fast runners, attaining speeds of up to 170 times their body length per second (Kamoun and Hogenhout, 1996). The analysis of morphological variation in this group is complicated due to low character variation and nondiscrete differentiation. We hence conducted comprehensive DNA sequencing across the geographic range of *Rivacindela* to provide the scaffold for assigning species membership. We find that the various existing procedures for DNA-based species delimitation identify biologically meaningful groups. The new likelihood method for analyzing branching rate produced results closely similar to those obtained with character-based methods but does not require prior assumption of population boundaries and can provide a statistical measure of confidence to the assessment of species limits.

MATERIAL AND METHODS

Collecting Information and DNA Procedures

Specimens were sampled at the edges of temporal salt lakes and salt flats in the Provinces of Western Australia, South Australia, and the Northern Territory during two expeditions in 2001 and 2003. Several of these sites were surveyed for cicindelids for the first time, and collections produced many apparently undescribed species and variants (Fig. 1). Specimens were placed directly into absolute ethanol after cursory identification in the field. Up to six individuals (more in a few cases) from each locality were sequenced, plus an additional set of up to six individuals if obvious morphologically different forms were recognized during field work at a given site or where individuals were obtained in different years or at multiple adjacent sites on the larger lakes (Supplementary Table S1; <http://systematicbiology.org>). This sampling regime was thought to be a good reflection of the total diversity encountered at the various collecting sites capturing species-level groups and their internal variation, given the absence of a thorough taxonomic treatment, while keeping the sequencing effort to a minimum. In total, 468 specimens assigned to 108 local sets of individuals (65 sites, plus morphologically distinct forms and samples from repeat visits of a site) were included in the analysis.

Nondestructive DNA extraction was performed using a Qiagen DNeasy kit. Whole specimens were soaked overnight in extraction buffer at 37°C, with small perforations made to the side of the abdomen and DNA




FIGURE 1. Map of 65 collecting sites. The distribution of four widespread species is shown as different symbols (species 1, light grey circles; species 11, black triangles; species 29, white boxes; species 35, dark grey diamonds). Detailed collecting information is provided in Supplementary Table S1.

extracted from the supernatant. Damage to specimens (some of them to be holotypes in future species descriptions) was minimal, and softness of specimens was maintained if transferred to 70% ethanol after the extraction. Vouchers will be maintained as pinned dry specimens in the W. D. Sumlin personal collection. Amplification of three mtDNA regions from the cytochrome oxidase subunit 1 (*cox1*), cytochrome *b* apoenzyme (*cob*), and 16S ribosomal RNA (*rrnL*) plus adjacent regions, was generally successful with well-established oligonucleotides (Pons et al., 2004) but required newly designed *cox1* and *rrnL* primers in some cases, the latter to avoid coamplification of a nuclear paralog (Pons and Vogler, 2005). (The genetic nomenclature in this paper follows Boore, 2001). Primer combinations for each region are given in Supplementary Table S2. PCR fragments were sequenced in both directions on an ABI377 automated sequencer. The sequences reported in this paper have been deposited in GenBank under accession nos. AJ617921–AJ618351 (*cox1*), AJ618352–AJ618766 (*cob*), AJ619087–AJ619548 (*rrnL*). In addition, three species of Australian cicindelids established to be closely related to *Rivacindela* based on a wider survey of *Cicindela sensu lato* (unpublished) were used as outgroups (AJ831553–AJ831564). The data

and tree files have been uploaded at Treebase under SN2798-10988.

Data Analysis

Statistical parsimony analysis was carried out with TCS version 1.3 software (Clement et al., 2000) using only individuals with complete sequence information for all three genes. Where a separate lake sample did not include at least one specimen with complete sequence information (listed in parentheses in Table S1), these were assigned to the networks based on their phylogenetic position in standard parsimony analysis (below). Statistical parsimony analysis partitions the data into independent networks of haplotypes connected by changes that are non-homoplastic with a 95% probability (Templeton, 2001). Intragroup and pairwise intergroup divergences were calculated with Mega version 2.1 (<http://www.megasoftware.net>) using p-distances. F_{st} and P values at 0.05 significance were calculated from pairwise differences in Arlequin 2.0 (<http://lgb.unige.ch/arlequin>).

Parsimony tree searches on the complete data set were conducted using PAUP*4.0b10 (Swofford, 2002) with


40 independent runs and 200 ratchet iterations (Nixon, 1999) each, with 15% of characters reweighted using the default settings of the PAUPRat script (<http://www.ucalgary.ca/~dsikes/software2.htm>). The preferred topology was selected among an initial set of parsimony trees, from which one was selected according to the Shimodaira-Hasegawa (SH) test under a GTR+I+ Γ likelihood model and used as starting tree for an ML search in PAUP*4.0b10. The ML search was conducted on a desktop computer for 3 days testing a total of 18,622 rearrangements. Where nodes were consistent with the ML tree, bootstrap values were given based on parsimony analysis obtained by searching 100 pseudoreplicated data sets (generated using SEQBOOT in Phylip 3.57) using the ratchet. A strict consensus tree was created from these trees for each pseudoreplicate, and the 100 strict consensus trees each representing a bootstrap replicate were used to create a 50% majority rule consensus tree providing the support values. Relative ages of nodes were estimated using the r8s software (Sanderson, 2002) by fitting branch lengths of an ML tree using penalized likelihood and a smoothing parameter of 10, chosen as optimal by cross-validation. Absolute ages were estimated setting the split of *R. aurifodina* (sample 204) and *R. salicursoria* (sample 208) to 3.2 Mya (Pons et al., 2004). This was based on an estimate from a phylogeny of worldwide lineages of *Cicindela sensu lato* calibrated from biogeographic evidence in North American lineages which included only these two species to represent *Rivacindela* (Barraclough and Vogler, 2002). Sequence divergence between individuals of these species was 6.0% (SD 0.14%) and hence the estimate corresponds closely to the widely used insect mtDNA molecular clock calibration of 2.3% per My (Brower, 1994). Lineage-through-time analyses were obtained as a semilogarithmic plot of the number of lineages against time since the first bifurcation (Nee et al., 1992).

Species Delimitation

Different quantitative methods for species delimitation were applied, implemented by visual inspection of the variable nucleotide positions and trees derived from these. These procedures were conducted only within the independent networks defined in the statistical parsimony analysis, greatly reducing the complexity of scoring separated groups. Population profiles of character variation were established according to Sites and Marshall (2003) as the basis for PAA (Davis and Nixon, 1992) and CHA (Brower, 1999), the latter by assessing variable characters on the likelihood tree shown below. The WP method also used this tree to delimit "exclusive" populations, defined as the monophyly of geographically restricted genotypes to the exclusion of clades elsewhere. F_{st} values were used for aggregating samples with non-significant pairwise F_{st} , in analogy to the grouping of populations under PAA (Supplementary Fig. S1).

Analyses of mtDNA Branching Times

A statistical model was developed to test for the predicted change in branching rates at the species boundary.



Waiting intervals	x1	x2	x3	x4	x5	etc.
Diversification (n)	2	3	2	2	1	etc.
Species 1 (n)	0	0	2	3	3	etc.
Species 2 (n)	0	0	0	0	0	etc.
Species 3 (n)	0	0	0	0	2	etc.

FIGURE 2. Schematic illustration of the waiting times in a calibrated tree and the numbers of lineages present for each type of diversification process (interspecies diversification and within-species coalescence) during each waiting interval. Branches are categorized as either between species (thin lines) or within species branching (bold lines) according to the procedures described in Material and Methods.

The overall aim of the procedure is to classify the observed branching time intervals defined by the nodes in a clock-constrained phylogram to either being the result of inter-specific ("diversification") or intraspecific ("coalescent") processes of lineage branching (Fig. 2). A full description of the model and its performance on simulated trees will be provided elsewhere (Barraclough, unpublished).

As raw data, the model uses the waiting times between successive branching events on the ultrametric DNA tree, x_i . We combine standard models that separately consider branching within populations (Hudson, 1991; Wakeley, 2006) and branching between species (Nee, 1994, 2001; Nee et al., 1994). Under a neutral coalescent, the likelihoods of the waiting times within a single population with effective population size N_e and n_i lineages present during waiting time i are given by:

$$L(x_i) = \lambda n_i (n_i - 1) e^{-\lambda n_i (n_i - 1) x_i} \quad (1)$$

where the birth rate

$$\lambda = \frac{1}{2N_e} \quad (2)$$

The simplest standard approach for considering branching between species is as a Yule model (Yule, 1924); i.e., a stochastic birth-only model. The likelihoods of the waiting times in a species phylogeny (one tip per species) of a clade with constant average speciation rate, λ , and no extinction are given by:

$$L_{(x_i)} = \lambda n_i e^{-\lambda n_i x_i} \quad (3)$$

We combine the above equations describing population and speciation processes to consider a clade that has diversified into k species, each of which can be treated as a single population with effective size, N_j , $j = 1$ to k . Assuming a constant speciation rate without extinction, and neutral coalescence within each species, the likelihoods of waiting times in the entire tree under this mixed Yule coalescent (MYC) model are given by:

$$L_{(x_i)} = b e^{-b x_i} \quad (4)$$

where

$$b = \lambda_{k+1} n_{i,k+1} + \sum_{j=1,k} (\lambda_j n_{i,j} (n_{i,j} - 1)) \quad (5)$$

where $k + 1$ is the index assigned to the diversification process and $n_{i,j}$ is the number of lineages in waiting interval i belonging to process j . λ_{k+1} is the speciation rate and λ_j are the branching rates for each coalescent process as defined in Equation 2. We do not consider effective population sizes explicitly hereafter, rather the coalescent branching rate parameter for each species. The term b is the probability that an event of any type happens at the end of the waiting interval and $e^{-b x_i}$ is the probability that no event happened during the waiting interval (see appendix 1 of Nee, 2001).

This MYC model can be fitted to an ultrametric tree by maximizing the sum of the log-likelihoods of waiting times across the entire tree. A key step is fitting the location of the switches from speciation to coalescent nodes; i.e., the most recent common ancestral node defining each species. The simplest approach is to assume that there is a threshold time, T , before which all nodes reflect diversification events and after which all nodes reflect coalescent events. Species in this model are thus delimited by the descendent nodes of branches crossing the threshold. T can be optimized to find the maximum likelihood solution and hence to estimate the number of species. Note that this approach does not assume that all species have the same age of their most recent common ancestor, which would not be the expected even under an equal population size model, but rather that the most recent diversification event occurred before the oldest within-species coalescent event. Approximate 95% confidence intervals for the parameters can be calculated by finding solutions with 2 log-likelihood units of the maximum (Edwards, 1972).

To test whether there is significant evidence for the predicted transition in branching rates, the likelihood

for the threshold model can be compared to that obtained assuming no threshold; i.e., a single branching process for the entire tree ($k = 0$). Assuming initially that all species have the same effective population size, the threshold version of the MYC model introduces two additional parameters compared to the null model: an additional branching rate parameter, λ_2 , and T . A standard log-likelihood ratio test can be used to assess whether the alternative model provides a significantly better fit than the null model of no such shift in branching process: twice the difference in log-likelihood is expected to be chi-square distributed with 2 degrees of freedom (Goldman, 1993). Failure to reject the null model could have several explanations. First, the clade might in fact represent a single species (unlikely in the present case). Second, the observed branching rate within species depends on the number of individuals sampled per species: small samples per species will weaken the power to detect the transition. Conversely, incomplete sampling at the species level would reduce the apparent branching rate in λ_1 . Third, some combinations of actual branching processes will also make it harder to detect the transition; for example, a combination of fast speciation rate and large population sizes.

Equations (4) to (6) make strict assumptions about both speciation and population processes that seem unlikely to hold in many real clades. As a general solution to this problem, we introduce additional parameters that allow for a qualitatively wide range of different models for both types of branching processes. Nee et al. (1994a) and Nee (2001) discussed transformations of waiting intervals that allow derived models to be treated as exponential processes with a single rate parameter. These accounts imply a simple general transformation of the MYC model by substituting b in Eq. (4) with b^* :

$$b^* = \lambda_{k+1} (n_{i,k+1})^{p_{k+1}} + \sum_{j=1,k} (\lambda_j (n_{i,j} (n_{i,j} - 1))^{p_j}) \quad (6)$$

We call this the general mixed Yule coalescent (GMYC) model. The p_j represent scaling parameters that can be optimized during model fitting. Interpretation depends on which class of branching events are considered. The scaling parameter for the diversification process, p_{k+1} , provides similar information to the gamma statistic of Pybus and Harvey (2000). $p_{k+1} = 1$ represents a constant speciation rate model with no extinction. $p_{k+1} > 1$ indicates an apparent increase in diversification rate towards the present, which might reflect, for example, a real increase in speciation rate or the effects of constant background extinction (Barraclough and Nee, 2001; Nee, 1994). $p_{k+1} < 1$ represents an apparent decrease in diversification rate towards the present, which might reflect, for example, a real slow-down in speciation or the effects of incomplete sampling of species within the clade (Nee et al., 1994b; Pybus and Harvey, 2000). The scaling parameters for the coalescent processes within each species are interpreted differently. $p_j = 1$ represents a neutral coalescent model. $p_j < 1$ indicates a relative deficit of

recent coalescent events, expected, for example, if populations were growing in size or experiencing balancing selection (Nee et al., 1994). $p_j > 1$ indicates a relative excess of recent coalescent events, expected, for example, if all populations were declining in size, following a recent selective sweep affecting the marker or if there is further population structure within the species. Hence, optimizing across possible values of p for both classes of branching events relaxes the assumptions of the method and can provide pointers for further analyses to explore possible causes for departure from the simplest model.

Code implementing the model in R using functions from the APE library is available from TGB. First, we visualized the waiting time data by plotting the log of the number of lineages through time (Nee et al., 1992). A transition in branching rates should be visible as a sudden increase in slope of the plot towards the present. Second, we ran the threshold version of the GMYC model on the calibrated ultrametric tree of *Rivacindela*. Third, we compared the likelihood to that obtained assuming a single branching process for the tree. For the GMYC model, assuming all species have the same parameter values, the threshold model has five parameters ($\lambda_1, \lambda_2, p_1, p_2$, and T), whereas the null model has two (λ_1 and p_1); hence, there are three degrees of freedom for the comparison.

RESULTS

Sequencing of three gene regions of mtDNA for the selected 468 specimens produced a data matrix of 1914 bps, of which 514 bps were informative. The average uncorrected p-distance between any two ingroup sequences was 5.8% (7.8%, 9.4%, and 2.8%, respectively, for the *cox1*, *cob*, and *rrnL* regions). We obtained 5355 parsimony trees of 3539 steps, and $CI = 0.280$ and $RI = 0.926$. Among those, one tree with highest likelihood was selected (from those which were significantly better in the SH test) as a starting tree in an ML search resulting in a slightly different topology and marginally improved likelihood score of $-\log L 22,363.56$ (Fig. 3). Basal relationships were characterized by relatively long internal branches, but tip branches were short and grouped clusters of very similar sequences, as would be expected if *Rivacindela* has split into isolated evolutionary entities, each consisting of closely related individuals.

Patterns of sequence variation were investigated for the presence of species-level groups based on various types of aggregation analysis (Fig. 4). An initial step of this analysis was to identify independent networks using statistical parsimony (Templeton et al., 1992), which separated the total variation into 25 groups based on connection limit of 15 steps; i.e., branches of 16 steps and beyond are considered to fall outside of the 95% confidence interval for these connections to be nonhomoplastic. The status of these independent networks as putative species was not questioned (see Sites and Marhall, 2003). Individual networks were then investigated for further subdivision. As we started the analysis with a total of 108 local samples whose coherence was not in doubt (i.e., they each constituted a set of specimens

collected at the same site and did not show any obvious morphological variation; Material and Methods), each of these were included in the various aggregation analyses conducted separately for each network. These analyses grouped the samples into either 46 (under PAA and CHA) or 47 (under WP analysis) separate entities, each entity consisting of up to 63 individuals from between 1 and 13 localities (Fig. 4). When analyzed separately, each of the three mtDNA regions still recovered 40 to 43 putative species of similar extent, and those obtained from 728 bp fragment of the *cox1* gene most closely matched the result from the full data set. Average pairwise divergence within these putative species determined by the WP method was 0.5%, much lower than the average among-species divergence of 6.3%, and 2.2% between sister species.

Population genetic analysis was also used for aggregation of populations with non-significant pairwise F_{st} (Supplementary Fig. S1), in analogy to PAA/CHA. Among the 66 groups with four or more individuals included in the analysis, under a significance level for F_{st} $P < 0.05$, this analysis led to recognition of only five additional entities over those in the WP analysis (Fig. 4). Bonferroni corrections for multiple comparisons would increase the significance level for F_{st} to $P = 0.00076$, which would lead to the recognition of fewer subdivided groups, although this value is likely to introduce a large type 2 error, i.e., the lumping of groups separated at a significant F_{st} level. Instead, using uncorrected pairwise F_{st} might result in type 1 error of accepting groups that appear separated simply due to chance, and therefore the reported results (Fig. 4) constitute the upper bound of the number of entities.

The existence of distinct mtDNA lineages was confirmed by analysis of branch lengths. A lineage-through-time plot showed a steep upturn in branching rates towards the present, presumably marking the transition from between-species to within-species rate of lineage branching (Fig. 5). The position of the transition was fitted using the GMYC model assuming a single threshold: pretransition nodes were assumed to reflect speciation events and posttransition nodes were assumed to reflect within species coalescence. The model fitted a transition in branching rate occurring at 0.43 Mya, leading to an estimate of 48 putative species, 5 of which contain a single individual. Confidence limits for the threshold ranged from 0.39 to 0.53 Mya and for the estimated number of species from 47 to 51. The GMYC model was preferred over the null model of uniform branching rates ($\log L = 2641$, compared to null model $\log L = 2606$; $2\Delta L = 71.8$, χ^2 test, d.f. = 3, $P \ll 0.001$). The groups of sequences identified by these shifts matched closely with those identified through other aggregation analyses (Fig. 4). There were six cases in which the WP method delimited species embedded within a paraphyletic species that were not detected by the branch-length method. Conversely there were seven cases of divergence (sometimes a single individual) detected by the branch-length analysis but not by the WP method.




FIGURE 3. ML tree depicting relationships of *Rivacindela* mtDNA haplotypes with branch lengths fitted assuming a molecular clock. The two main sister groups are shown separately (panel A, Eastern group; panel B, Western group). Localities for each of 47 putative species (“Clades”) obtained under the WP method are indicated by a two-letter code. Different shading is used for better visibility of the extent of groups. Gray bars and site names colours indicate widely distributed or paraphyletic species. The line style indicates whether branches were estimated as between-species branching (stippled) or within-species branching (solid) in the likelihood procedure. The dotted vertical line shows the maximum likelihood transition point of the switch in branching rates. The grey shading indicates the confidence limits for the transition point falling within 2 log-likelihood units of the ML solution (Material and Methods). Numbers above nodes represent bootstrap support values based on 100 pseudoreplicates and ratchet parsimony searches, shown only for the deep level clades. Bootstrap values under 50% are not shown. (Continued)




FIGURE 3. (Continued)

Most putative species identified in the analysis were restricted to a single lake, but four widely distributed species were present at between 5 and 13 sites, showing contiguous ranges that included all nearest neighbor sites (Fig. 1). Geographical structure was apparent also at deeper levels in the tree. For example, the basal node defined two major clades roughly coincident with eastern and western localities (Fig. 3). Sister species and larger sister clades tended to be found mostly at neighboring lakes or lake systems. A major divide separating the drainage systems (van de Graaff et al., 1977) was also apparent in the distribution of haplotypes. In western Australia, most populations of the same basin or paleo-river system were included in the same clade composed of closely related haplotypes. Haplotypes up to approximately 0.7 My of divergence were almost entirely confined to a single paleo-river, whereas populations from different drainage systems were not closely related, even where these lakes were in close geographic proximity (Fig. 6). Although most lakes harbored only one species, a maximum of three co-occurring species was observed, but these were not closely related in most cases.

DISCUSSION

The use of short mtDNA sequences in taxonomic studies, including "DNA barcoding" (Hebert et al., 2003), has been applied mostly to known species, but beyond novel tools for identification (and occasional evidence for cryptic species; Hebert et al., 2004a, 2004b; Meyer and Paulay, 2005), this does not directly bear on the taxonomic status of groups (Moritz and Cicero, 2004). Here we used quantitative analyses of sequence data to delimit putative species, whereby the sequence information itself provided a framework for alpha taxonomy. Species delimitation relied on two principal approaches, including established procedures based on diagnostic characters (Sites and Marshall, 2003) and a novel procedure based on detecting the shift in the rate of lineage branching. Other studies have observed similar shifts in branching rates as signatures of the species boundary (e.g., Acinas et al., 2004; Barraclough et al., 2003; Cardoso and Vogler, 2005; Hebert and Gregory, 2005; Hugall et al., 2002; Monaghan et al., 2005; Wiens and Penkrot, 2002). However, to our knowledge there have been no previous attempts to utilise this information in a quantitative procedure of species delimitation, despite the development of methods for clustering of sequences (Blaxter, 2004) and matching them to predefined taxonomic groups (Steinke et al., 2005; Matz and Nielsen, 2005).

The branch-length analysis relies on a probabilistic model separating species diversification (phylogeny) from coalescent processes (genealogy within species). In this sense, the shift in branching rate corresponds to the elusive boundary between Hennig's (1966) phylogenetic and tokogenetic realms, indicating divergent and reticulate relationships. Our goal was to produce a method capable of fitting approaches of phylogenetics and population genetics, as the two major disciplines operating above and below this boundary (Brower et al., 1996). In

its current implementation, the approach is based on simple assumptions such as a constant speciation rate model or neutral coalescence, although these are relaxed somewhat by incorporating the scaling parameter. The procedure has acceptable false-positive rates when applied to data simulated assuming no transition in branching rates and reasonable power for data simulated assuming distinct species (Barraclough, unpublished). The approach could be modified to specify detailed models of speciation, extinction, and population processes, and then to choose among competing models, but because of the bewildering range of possible scenarios (Barraclough and Nee, 2001; Charlesworth et al., 2003), this was not further explored here.

The analysis of branch length overcomes two principal problems of existing methods for quantitative species delimitation. First, the approach takes into account uncertainty of species limits by permitting confidence intervals when allocating species-defining nodes, a desirable property where species limits are weakly developed (Hey et al., 2003). Population aggregation methods do not allow for uncertainty, but rather optimize species limits based on a fixed criterion. Other recent population-based approaches take uncertainty into account (e.g., Matz and Nielsen, 2005) but have yet to be applied to species delimitation per se rather than identification of unknown sequences. Second, the branch length-based methods do not require that populations are defined a priori before being subjected to tests of aggregation (Davis and Nixon, 1992; Mayden, 1997). Definitions of population units remain problematic, as they frequently lack easily discernible spatial or genetic boundaries (Schaefer, 2006); e.g., where distributional ranges are insufficiently studied or where morphologically cryptic forms are found in the same geographical locality. For our study of *Rivacindela* we simplistically assumed that individuals collected at different sites were members of separate populations, whereas all individuals from a single site were considered a uniform population, unless morphological differences were evident. As population aggregation analyses are based on the a priori definitions of populations, this kind of informal treatment could introduce a major factor of uncertainty into species delimitation. The branching-based approach is independent of the recognition of population limits. In addition, the approach permits the inclusion of rare species represented by only a single individual, which is problematic in population based species concepts.

The length of time needed to attain the signatures used for species recognition differs between these methods. For example, CHA and PAA rely on the prediction that fixed nucleotide differences are unlikely between random samples of a single population, but expected to accumulate between isolated populations. Under a neutral model of mutation and genetic drift, for a mitochondrial marker the expected time to the first fixed difference, and hence ability of these methods to detect isolated species, is less than $0.5 Ne$ generations after populations were isolated (Hey, 1991). The WP method relies on the assumption that an isolated population will




FIGURE 6. Distribution of species in paleo-drainages for the western Australian assemblages (Panel B of tree shown in Fig. 3; see Fig. 1 for an outline map of the wider region). Shadings and symbols indicate different paleo-river systems according to fig. 5 in van de Graaff et al. (1977). The dashed lines mark lineages inferred to have undergone a shift between major drainage systems marked by the arrows in the map (inset).

eventually become monophyletic with respect to its ancestral or sister population. According to simulations of the probability to observe reciprocal monophyly, this is expected to occur at around $0.7 Ne$ generations; i.e., there

is a 50% chance of a given locus being monophyletic at that time (table 1 in Hudson and Coyne, 2002). The branch-length analysis proposed here requires additionally that monophyletic groups are recognizable on longer

stem branches, expected to occur around Ne generations after lineage separation (Hudson and Coyne, 2002). To have 95% of a given locus displaying a given signature, the time interval needed will be higher in each case, for example, over $2N_e$ generations, for changes in branching rate to arise. The final method we used, based on the inbreeding coefficient, detects significant between-population genetic subdivision from pairwise F_{st} estimates, which does not entail fixed differences or monophyly. Hence this method might detect more recent breaks in gene flow than the others, but risks that populations with partial gene flow are recognized as separate entities: it constitutes an extreme upper limit for species numbers in population-based analyses.

Whether these methods actually produce differing estimates of numbers of groups will depend on the relative separation of species versus within-species coalescence times. If many related species diverged more recently than the oldest coalescence times of the alleles they contain, the methods will underestimate species numbers, delimiting species complexes rather than individual species. This will affect methods relying on branch length more strongly than methods based on diagnostic fixed character states which have a greater chance to identify recently diverged or otherwise cryptic species because they are expected to be apparent sooner after divergence (Goldstein and Desalle, 2003). However, in the present case very similar estimates of species numbers and species limits were obtained with either method, demonstrating the robustness of the data to different quantitative approaches of species delimitation. Differences that did occur reflect the opposing effects of cryptic within-population divergence detected by the analysis of branching rates on the one hand, and the recognition of more recent divergence by population-based methods on the other. In addition, the analysis of branching rates conducted on monophyletic lineages will not recognize paraphyletic species, while also relying on the correct reconstruction of the tree. Arguably, the best final assignment would be to integrate branch-length approaches and the PAA method (as it is not based on topology and hence a "correct" tree) and to recognize 54 putative species.

Neither approach would recognize recently derived ecological variants. For example, *R. eburneola* is a flightless species from Lake Gilmore, which was not separated by any of the DNA methods as distinct from the co-occurring, but flighted and morphologically and ecologically distinct *R. nr. blackburni*. This form should clearly be added to the count of *Rivacindela* species. The example hence demonstrated that the mtDNA-based methods may be conservative compared to classical morphological techniques of species differentiation, contrary to widely held concerns (see Agapow et al., 2004). In addition, DNA delimited species were generally congruent when compared to the morphologically based species assignment in those cases where names had been assigned to the specimens in preliminary identifications (Table S1), which allays concerns that DNA-based taxonomies result in oversplitting of the existing variation.

Although *R. eburneola* was the only obvious case where major morphological differences did not coincide with the mtDNA defined groups, minor variation was seen in the informal morphological identification during field work that also disagreed with DNA (Table S1). This raises the possibility of incongruent gene and species histories, in particular in the light of reported high propensity for gene flow of mtDNA between otherwise separated groups (Funk and Omland, 2003; Hudson and Coyne, 2002; Seberg et al., 2003; Will and Rubinoff, 2004). Without further morphological analysis or sequencing of nuclear loci, the magnitude of this problem in *Rivacindela* is difficult to assess. However, the high fidelity of geographic distribution of DNA-based groups and ancient river basins (Fig. 6) provides clear evidence against mtDNA dispersal, in particular during the past ~700 Ky since most of these species were separated. Across all western Australian species we observe only two cases of well-sampled distinct clusters in the branch lengths analysis (individuals from site GY in clade 30, and individuals from site DE in clade 28; the DE individuals being a diagnosable group in the PAA) where one or more sequences were embedded in a "foreign" population; i.e., in a different river basin than all other sequences within the cluster. Hence, in *Rivacindela* the broad congruence of mtDNA distribution with biogeographical landscape features is evidence for their biological relevance.

The geographic pattern strongly indicates vicariant speciation as the cause of the range distributions, with only occasional dispersal between drainage systems, consistent with the paleoclimatic history of the region. Ephemeral salt lakes of interior Australia formed by fragmentation of paleo-drainage basins that existed prior to the aridification of the continent starting in the Miocene (van de Graaff et al., 1977). Proposed ancient basins remained well preserved because of tectonic stability and slow erosion and sedimentation in the area (van de Graaff et al., 1977). The date of the species-to-population transition (Fig. 4) coincides narrowly with the changes from lacustrine clays to evaporites and dune sediments estimated to have occurred between 400 and 700 Kya in a final shift to present-day aridity in regional climates (Pillans and Bourman, 2001). This would indicate that species formation in *Rivacindela* was a direct result of the fragmentation of their habitat near edges of the disappearing river systems.

CONCLUSIONS

The acceptance of DNA taxonomy in the literature has been slow (e.g., Lipscomb et al., 2003; Wheeler, 2004; Will and Rubinoff, 2004), partly because the biological status of entities established from sequence information alone is not easily verified, unlike the long tradition of taxonomic research based on morphological characters. However, in *Rivacindela* the species status of the mtDNA-based entities is well justified both evolutionarily and biologically: the levels of sequence divergence between species matched those in other insects (Brower, 1994), including taxonomically well studied North American

Cicindela (Vogler et al., 2005); geographic ranges are contiguous but clearly delimited and consistent with biogeography (Fig. 1); patterns of genetic diversity are consistent with paleoclimatic evidence for their vicariant origin (Fig. 6); and the tree exhibits the branching rate signature of the species-to-population transition (Fig. 5). Besides the question about species status, the mtDNA groups also provide a powerful system for taxonomic communication, identification, and convenient integration of newly generated information. For example, specimens found during our second (2003) expedition were readily identified in the existing database either as members of established species or representing unknown groups (Supplementary Table S1).

With an estimated 10^7 species of insects on Earth (Erwin, 1982; Odegaard, 2000), there is no realistic hope for their in-depth taxonomic treatment using traditional methods. Greater sequencing throughput and algorithmic analysis of species membership, as applied here to a depauperate desert ecosystem, could be extended to more complex assemblages, including the spectacular diversity of insect species in tropical forests. The efficacy of such approaches will depend on whether similar branching rate shifts are found generally in insects and to what degree they separate closely related species. The prominent feature of the branch-length distribution of *Rivacindela* defined by the largely synchronized origin of species fitting a simplified model of rate shift may be a special case due to its unique paleoclimatic history. In addition, the extent of population sampling in a large clade may rarely be as complete as in the current study, and could also affect the recovery of the rate shift, suppressing or inflating rates in either portion of the tree and shifting their slope relative to each other. Finally, population subdivision within species would also blur the species-to-population transition of branching rates, as subdivision (metapopulation structure) tends to increase the age of the most recent common ancestor within species (Wakeley, 2000; Wakeley and Aliacar, 2001), although, fortuitously, if the common ancestor of diverged species was also subdivided, genetic divergence between sister species will tend to increase by a similar amount (Wakeley, 2000). Given the emerging evidence for strong clustering of sequence variation in most groups of insects and other invertebrates (Hajibabaei et al., 2006; Meyer and Paulay, 2005; Monaghan et al., 2005; Page et al., 2005; Smith et al., 2005) and their largely uniform coalescence time (Meyer and Paulay, 2005), for the great majority of lineages the signal of the clade-to-population transition is probably sufficiently robust to be detectable even under a range of confounding biological factors and sampling regimes. This supports the wide applicability of mtDNA-based species delimitation in poorly known groups generally.

As well as establishing a taxonomic framework, the high-throughput mtDNA profiles simultaneously provide information on the evolutionary and ecological history of a group. In *Rivacindela*, we established that macroecological patterns (endemism, beta-diversity,

range sizes) and dynamics of speciation result largely from the history of climate change, whereby separated populations have persisted in situ for 10^5 to 10^6 years. This provides a context for comparative studies, e.g., with the subterranean diving beetles dependent on the same aquifers as *Rivacindela* (Leys et al., 2003), or with patterns of diversity in the 1000 species of *Cicindela s.l.* on other continents (Pearson and Vogler, 2001). While providing a framework for taxonomy widely accessible without specialist knowledge, the comprehensive mtDNA profiling of faunas across biogeographic regions (see also Monaghan et al., 2006) at the same time provides a formidable resource for evolutionary biology, population genetics, and studies of global biodiversity patterns.

ACKNOWLEDGMENTS

We thank David Brzoska, Fabio Cassola, and Jay Shetterly for help with collection of specimens, and Saskia Bode and Fatos Kopliku for sequencing. Charles Godfray, Ryan Gregory, Paul Harvey, Andy Purvis, Sean Nee, and Richard Thomas provided helpful comments on an earlier version of the manuscript. We are grateful to Marshal Hedin, Rod Page, Mark Fishbein, and an anonymous referee for their detailed and insightful comments in the review process. A permit for export of specimens from Australia was held by WDS. This work was funded by the Natural Environment Research Council (UK) grant NER/A/S/2000/00489 and NER/A/S/2001/01133.

REFERENCES

- Acinas, S. G., V. Klepac-Ceraj, D. E. Hunt, C. Pharino, I. Ceraj, D. L. Distel, and M. F. Polz. 2004. Fine-scale phylogenetic architecture of a complex bacterial community. *Nature* 430:551–554.
- Agapow, P. M., O. R. P. Bininda-Emonds, K. A. Crandall, J. L. Gittleman, G. M. Mace, J. C. Marshall, and A. Purvis. 2004. The impact of species concept on biodiversity studies. *Quart. Rev. Biol.* 79:161–179.
- Barracough, T. G., C. W. Birky, and A. Burt. 2003. Diversification in sexual and asexual organisms. *Evolution* 57:2166–2172.
- Barracough, T. G., and S. Nee. 2001. Phylogenetics and speciation. *Trends Ecol. Evol.* 16:391–399.
- Barracough, T. G., and A. P. Vogler. 2002. Recent diversification rates in North American tiger beetles (genus *Cicindela*). *Mol. Biol. Evol.* 19:1706–1716.
- Blaxter, M. L. 2004. The promise of a DNA taxonomy. *Phil. Trans. R. Soc. B* 359:669–679.
- Boore, J. L. 2001. Mitochondrial gene arrangement source guide, version 6.0. http://evogen.jgi.doe.gov/second.levels/mito-chondria/MGA_Guide.html. Department of the Environment, Joint Genome Institute, Walnut Creek, California.
- Brower, A. V. Z. 1994. Rapid morphological radiation and convergence among races of the butterfly *Heliconius erato* Inferred from patterns of mitochondrial DNA Evolution. *Proc. Natl. Acad. Sci. USA* 91:6491–6495.
- Brower, A. V. Z. 1999. Delimitation of phylogenetic species with DNA sequences: A critique of Davis and Nixon's population aggregation analysis. *Syst. Biol.* 48:199–213.
- Brower, A. V. Z., R. DeSalle, and A. Vogler. 1996. Gene trees, species trees, and systematics. *Annu. Rev. Ecol. Syst.* 27:423–450.
- Cardoso, A., and A. P. Vogler. 2005. DNA taxonomy, phylogeny and Pleistocene diversification of the *Cicindela hybrida* species group (Coleoptera: Cicindelidae). *Mol. Ecol.* 14:3531–3546.
- Charlesworth, B., D. Charlesworth, and N. H. Barton. 2003. The effects of genetic and geographic structure on neutral variation. *Annu. Rev. Ecol. Syst.* 34:99–125.
- Clement, M., D. Posada, and K. A. Crandall. 2000. TCS: A computer program to estimate gene genealogies. *Mol. Ecol.* 9:1657–1659.
- Cracraft, J. 1983. Species concept and speciation analysis. *Curr. Ornith.* 1:159–187.

- Cracraft, J. 1992. The species of the birds-of-paradise (Paradisaeidae): Applying the phylogenetic species concept to complex pattern of diversification. *Cladistics* 8:1–43.
- Davis, J. I., and K. C. Nixon. 1992. Populations, genetic variation, and the delimitation of phylogenetic species. *Syst. Biol.* 41:421–435.
- Doyle, J. J. 1995. The irrelevance of allele tree topologies for species delimitation, and a non-topological alternative. *Syst. Bot.* 20:574–588.
- Edwards, A. W. F. 1972. *Likelihood*. The John Hopkins University Press, London.
- Erwin, T. L. 1982. Tropical forests: Their richness in Coleoptera and other arthropod species. *Coleopt. Bull.* 36:74–75.
- Freitag, R. 1979. Reclassification, phylogeny, and zoogeography of the Australian species of *Cicindela* (Coleoptera: Cicindelidae). *Aust. J. Zool. Suppl. Ser.* 66:1–99.
- Funk, D. J., and K. E. Omland. 2003. Species-level paraphyly and polyphyly: Frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annu. Rev. Ecol. Syst.* 34:97–423.
- Goldman, N. 1993. Statistical tests of models of DNA substitution. *J. Mol. Evol.* 36:182–198.
- Goldstein, P. Z., and R. Desalle. 2003. Calibrating phylogenetic species formation in a threatened insect using DNA from historical specimens. *Mol. Ecol.* 12:1993–1998.
- Hajibabaei, M., D. H. Janzen, J. M. Burns, W. Hallwachs, and P. D. N. Hebert. 2006. DNA barcodes distinguish species of tropical Lepidoptera. *Proc. Natl. Acad. Sci. USA* 103:968–971.
- Hebert, P. D. N., A. Cywinska, S. L. Ball, and J. R. DeWaard. 2003. Biological identifications through DNA barcodes. *Proc. R. Soc. B* 270:313–321.
- Hebert, P. D. N., and T. R. Gregory. 2005. The promise of DNA barcoding for taxonomy. *Syst. Biol.* 54:852–859.
- Hebert, P. D. N., E. H. Penton, J. M. Burns, D. H. Janzen, and W. Hallwachs. 2004a. Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptus fulgurator*. *Proc. Natl. Acad. Sci. USA* 101:14812–14817.
- Hebert, P. D. N., M. Y. Stoeckle, T. S. Zemlak, and C. M. Francis. 2004b. Identification of birds through DNA barcodes. *PLoS Biol.* 2:1657–1663.
- Hennig, W. 1966. *Phylogenetic systematics*. University of Illinois Press, Urbana.
- Hey, J. 1991. The structure of genealogies and the distribution of fixed differences between DNA-sequence samples from natural populations. *Genetics* 128:831–840.
- Hey, J., R. S. Waples, M. L. Arnold, R. K. Butlin, and R. G. Harrison. 2003. Understanding and confronting species uncertainty in biology and conservation. *Trends Ecol. Evol.* 18:597–603.
- Hudson, R. R. 1991. Gene genealogies and the coalescent process. *Oxf. Surv. Evol. Biol.* 7:1–44.
- Hudson, R. R., and J. A. Coyne. 2002. Mathematical consequences of the genealogical species concept. *Evolution* 56:1557–1565.
- Hugall, A., C. Moritz, A. Moussalli, and J. Stanisc. 2002. Reconciling paleodistribution models and comparative phylogeography in the Wet Tropics rainforest land snail *Gnarosophia bellendenkerensis* (Brazier 1875). *Proc. Natl. Acad. Sci. USA* 99:6112–6117.
- Kamoun, S., and S. A. Hogenhout. 1996. Flightlessness and rapid terrestrial locomotion in tiger beetles of *Cicindela* L. subgenus *Rivacindela* van Nidek from saline habitats of Australia. *Coleopt. Bull.* 50:221–230.
- Leys, R., C. H. S. Watts, S. J. B. Cooper, and W. F. Humphreys. 2003. Evolution of subterranean diving beetles (Coleoptera: Dytiscidae: Hydroporini, Bidessini) in the arid zone of Australia. *Evolution* 57:2819–2834.
- Lipscomb, D., N. Platnick, and Q. Wheeler. 2003. The intellectual content of taxonomy: A comment on DNA taxonomy. *Trends Ecol. Evol.* 18:65–68.
- Mallet, J., and K. Willmott. 2003. Taxonomy: Renaissance or Tower of Babel? *Trends Ecol. Evol.* 18:57–59.
- Matz, M. V., and R. Nielsen. 2005. A likelihood ratio test for species membership based on DNA sequence data. *Phil. Trans. R. Soc. B* 360:1969–1974.
- Mayden, R. L. 1997. A hierarchy of species concepts: The denouement in the saga of the species problem. Pages 381–423 in *Species. The units of biodiversity* (M. F. Claridge, H. A. Dawah, and M. R. Wilson, eds.). Chapman and Hall, London.
- Meyer, C. P., and G. Paulay. 2005. DNA barcoding: Error rates based on comprehensive sampling. *PLoS Biol.* 3:2229–2238.
- Monaghan, M. T., M. Balke, T. R. Gregory, and A. P. Vogler. 2005. DNA-based species delineation in tropical beetles using nuclear and mitochondrial markers. *Phil. Trans. R. Soc. B* 360:1925–1933.
- Monaghan, M. T., M. Balke, J. Pons, and A. P. Vogler. 2006. Beyond barcodes: complex DNA taxonomy of a South Pacific Island radiation. *Proc. R. Soc. B* 273:887–893.
- Moritz, C., and C. Cicero. 2004. DNA barcoding: Promise and pitfalls. *PLoS Biol.* 2:1529–1531.
- Nee, S. 1994. Extinction rates can be estimated from molecular phylogenies. *Phil. Trans. R. Soc. B* 344:77–82.
- Nee, S. 2001. Inferring speciation rates from phylogenies. *Evolution* 55:661–668.
- Nee, S., R. M. May, and P. H. Harvey. 1994. The reconstructed evolutionary process. *Phil. Trans. R. Soc. B* 344:305–311.
- Nee, S., A. O. Mooers, and P. H. Harvey. 1992. Tempo and mode of evolution revealed from molecular phylogenies. *Proc. Natl. Acad. Sci. USA* 89:8322–8326.
- Nelson, G. J., and N. I. Platnick. 1981. *Systematics and biogeography: Cladistics and vicariance*. Columbia University Press, New York.
- Nixon, K. C. 1999. The Parsimony Ratchet, a new method for rapid parsimony analysis. *Cladistics* 15:407–414.
- Odegaard, F. 2000. How many species of arthropods? Erwin's estimate revised. *Biol. J. Linn. Soc.* 71:583–597.
- Page, T. J., S. C. Choy, and J. M. Hughes. 2005. The taxonomic feedback loop: Symbiosis of morphology and molecules. *Biol. Lett.* 1:139–142.
- Pearson, D. L., and A. P. Vogler. 2001. *Tiger beetles: The ecology and evolution of the Cicindelidae*. Cornell University Press, Ithaca, New York.
- Pillans, B., and R. Bourman. 2001. Mid Pleistocene arid shift in southern Australia, dated by magnetostratigraphy. *Aust. J. Soil Res.* 39:89–98.
- Pons, J., T. G. Barraclough, K. Theodorides, A. Cardoso, and A. P. Vogler. 2004. Using exon and intron sequences of the gene *Mp20* to resolve basal relationships in *Cicindela* (Coleoptera:Cicindelidae). *Syst. Biol.* 53:554–570.
- Pons, J., and A. P. Vogler. 2005. Complex pattern of coalescence and fast evolution of a mitochondrial rRNA pseudogene in a recent radiation of tiger beetles. *Mol. Biol. Evol.* 22:991–1000.
- Rosenberg, N. A., and M. Nordborg. 2002. Genealogical trees, coalescent theory and the analysis of genetic polymorphisms. *Nat. Rev. Genet.* 3:380–390.
- Sanderson, M. J. 2002. Estimating absolute rates of molecular evolution and divergence times: A penalized likelihood approach. *Mol. Biol. Evol.* 19:101–109.
- Schaefer, J. A. 2006. Towards maturation of the population concept. *Oikos* 112:236–240.
- Seberg, O., C. J. Humphries, S. Knapp, D. W. Stevenson, G. Petersen, N. Scharff, and N. M. Andersen. 2003. Shortcuts in systematics? A commentary on DNA-based taxonomy. *Trends Ecol. Evol.* 18:63–65.
- Sites, J. W., and J. C. Marshall. 2003. Delimiting species: A Renaissance issue in systematic biology. *Trends Ecol. Evol.* 18:462–470.
- Smith, M. A., B. L. Fisher, and P. D. N. Hebert. 2005. DNA barcoding for effective biodiversity assessment of a hyperdiverse arthropod group: the ants of Madagascar. *Phil. Trans. R. Soc. B* 360:1825–1834.
- Steinke, D., M. Vences, W. Salzburger, and A. Meyer. 2005. TaxI: A software tool for DNA barcoding using distance methods. *Phil. Trans. R. Soc. B* 360:1975–1980.
- Sumlin, W. D. 1997. Studies on the Australian Cicindelidae XII: Additions to *Megacephala*, *Nickerlea* and *Cicindela* with notes (Coleoptera). *Bull. Worldw. Res.* 4:1–56.
- Swofford, D. L. 2002. *PAUP*: Phylogenetic analysis using parsimony. Version 4.0b*. Sinauer Associates, Sunderland, Massachusetts.
- Tautz, D., P. Arctander, A. Minelli, R. H. Thomas, and A. P. Vogler. 2003. A plea for DNA taxonomy. *Trends Ecol. Evol.* 18:70–74.
- Templeton, A. R. 2001. Using phylogeographic analyses of gene trees to test species status and processes. *Mol. Ecol.* 10:779–791.
- Templeton, A. R., K. A. Crandall, and C. F. Sing. 1992. A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and sequencing data. III. Cladogram estimation. *Genetics* 132:619–633.

- Templeton, A. R., E. Routman, and C. A. Phillips. 1995. Separating population structure from population history: a cladistic analysis of the geographical distribution of mitochondrial DNA haplotypes in the tiger salamander, *Ambystoma tigrinum*. *Genetics* 140:767–782.
- van de Graaff, W. J. E., R. W. A. Crowe, J. A. Bunting, and M. J. Jackson. 1977. Relict early Cainozoic drainages in arid Western Australia. *Zeitschrift f. Geomorphol.* 21:379–400.
- Vogler, A. P., A. Cardoso, and T. G. Barraclough. 2005. Exploring rate variation among and within sites in a densely sampled tree: Species level phylogenetics of North American tiger beetles (Genus *Cicindela*). *Syst. Biol.* 54:4–20.
- Vrana, P., and W. C. Wheeler. 1992. Individual organisms as terminal entities: Laying the species problem to rest. *Cladistics* 8:67–72.
- Wakeley, J. 2000. The effects of subdivision on the genetic divergence of populations and species. *Evolution* 54:1092–1101.
- Wakeley, J. 2006. *Coalescent theory: An introduction*. Roberts & Co., Greenwood Village, Colorado.
- Wakeley, J., and N. Aliacar. 2001. Gene genealogies in a metapopulation. *Genetics* 159:893–905.
- Wheeler, Q. D. 2004. Taxonomic triage and the poverty of phylogeny. *Phil. Trans. R. Soc. B* 359:571–583.
- Wiens, J. J., and T. A. Penkrot. 2002. Delimiting species using DNA and morphological variation and discordant species limits in spiny lizards (*Sceloporus*). *Syst. Biol.* 51:69–91.
- Wilder, J. A., and H. Hollocher. 2003. Recent radiation of endemic Caribbean *Drosophila* of the dunni subgroup inferred from multilocus DNA sequence variation. *Evolution* 57:2566–2579.
- Will, K. W., and D. Rubinoff. 2004. Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics* 20:47–55.
- Yule, G. U. 1924. A mathematical theory of evolution based on the conclusions of Dr. J. C. Willis, FRS. *Phil. Trans. R. Soc. B* 213:21–87.

First submitted 16 October 2005; reviews returned 31 December 2005;

final acceptance 7 April 2006

Associate Editor: Marshal Hedin